

# DETECTING AND TREATING DEPRESSION IN ADULTS

## REFERENCES

1. Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR et al. The epidemiology of major depressive disorder: results from the national comorbidity survey replication (NCS-R). *JAMA*. 2003;289:3095-3105.
2. Hasin DS, Goodwin RD, Stinson FS, Grant BF. Epidemiology of major depressive disorder: results from the national epidemiologic survey on alcoholism and related conditions. *Arch Gen Psychiatry*. 2005;62(10):1097-1106.
3. NYC DOHMH, Press Release, March 12, 2007. Available at: [www.nyc.gov/html/doh/html/pr2007/pr014-07.shtml](http://www.nyc.gov/html/doh/html/pr2007/pr014-07.shtml).
4. Horowitz JA, Goodman JH. Identifying and treating postpartum depression. *J Obstet Gynecol Neonatal Nurs*. 2005;32(2):264-273.
5. Gaynes BN, Gavin N, Meltzer-Brody S, Lohr KN, Swinson T, Gartlehner G, et al. Perinatal depression: prevalence, screening accuracy, and screening outcomes. *Evid Rep Technol Assess (Summ)*. 2005;119:1-8.
6. Kim-Cohen J, Moffitt TE, Taylor A, Pawlby SJ, Caspi A. Maternal depression and children's antisocial behavior: nature and nurture effects. *Arch Gen Psychiatry*. 2005;62(2):173-18.
7. Lieb, R., Isensee, B., Hofler, M., Pfister, H., et al. Parental major depression and the risk of depression and other mental disorders in offspring: a prospective longitudinal study. *Arch Gen Psychiatry*. 2002;59(4):365-374.
8. Pilowsky DJ, Wickramaratne PJ, Rush AJ, Hughes CW, Garber J, Malloy E, King CA, Cerdá G, Sood AB, Alpert JE, Wisniewski SR, Trivedi MH, Talati A, Carlson MM, Liu HH, Fava M, Weissman MM. Children of currently depressed mothers: a STAR\* D ancillary study. *J Clin Psychiatry*. 2006;67(1):126-136.
9. Ballenger JC, Davidson JR, Leclerc Y, et al. Consensus statement on the primary care management of depression from the International Consensus Group on Depression and Anxiety. *J Clin Psychiatry*. 1999;60(suppl 7):54-61.
10. Conwell Y. Suicide in elderly patients. In: Schneider LS, Reynolds CF III, Lebowitz BD, Friedhoff A, eds. *Diagnosis and treatment of depression in late life: results of the NIH Consensus Development Conference*. Washington, DC: American Psychiatric Publishing;1994:397-418.
11. Simon GE, VonKorff M, Piccinelli M, Fullerton C, Ormel J. An international study of the relation between somatic symptoms and depression. *N Engl J Med*. 1999;341(18):1329-1335.
12. Co-Occurring Major Depressive Episode (MDE) and Alcohol Use Disorder Among Adults. *The National Survey on Drug Use and Health Report*. February 16, 2007; full report available at: [www.oas.samhsa.gov](http://www.oas.samhsa.gov).
13. Zimmerman M, McDermut W, Mattia J. Frequency of anxiety disorders in psychiatric outpatients with major depressive disorder. *Am J Psychiatry*. 2000;157(8):1337-1340.
14. Lasser K, Boyd JW, Woolhandler S, Himmelstein DU, McCormick D, Bor DH. Smoking and mental illness: a population-based prevalence study. *JAMA*. 2000;284(20):2606-2610.
15. Grant BF, Hasin DS, Chou SP, Stinson FS, Dawson DA. Nicotine dependence and psychiatry disorders in the United States: results from the national epidemiology survey on alcohol and related conditions. *Arch Gen Psychiatry*. 2004;61(11):1107-1115.
16. Lesperance F, Frasure-Smith N, Talajic M. Major depression before and after myocardial infarction: its nature and consequences. *Psychosom Med*. 1996;58(2):99-110.
17. Anderson RJ, Freedland KE, Clouse RE, Lustman PJ. The prevalence of comorbid depression in adults with diabetes: a meta-analysis. *Diabetes Care*. 2001;24(6):1069-1078.
18. Pohjasvuo T, Leppävuori A, Siira I, Vataja R, Kaste M, Erkinjuntti T. Frequency and clinical determinants of poststroke depression. *Stroke*. 1998;29(11):2311-2317.
19. Hackett ML, Anderson CS. Predictors of depression after stroke: a systematic review of observational studies. *Stroke*. 2005;36(10):2296-2301.
20. Starkstein S, Jorge R, Mizrahi R, Robinson R. The construct of minor and major depression in Alzheimer's Disease. *Am J Psychiatry*. 2005;162(11):2086-2093.
21. Bing EG, Burman MA, Longshore D, Fleishman JA, Sherbourne CD, London AS, et al. Psychiatric disorders and drug use among human immunodeficiency virus-infected adults in the United States. *Arch Gen Psychiatry*. 2001;58(8):721-728.
22. Whooley MA. Case-finding instruments for depression. *J Gen Intern Med*. 1997;12(7):439-445.
23. Arroll B, Khin N, Kerse N. Screening for depression in primary care with two verbally asked questions: cross sectional study. *BMJ*. 2003;327(7424):1144-1146.
24. Kroenke K, Spitzer RL, Williams JB. The PHQ-9. Validity of a brief depression severity measure. *J Gen Intern Med*. 2001;16(9):606-613.
25. Practice guideline for the treatment of patients with major depressive disorder (revision). American Psychiatric Association. *Am J Psychiatry*. 2000;157(4 Suppl):1-45.
26. American Psychiatric Association. Practice guideline for the treatment of patients with bipolar disorder. Available at: [www.psych.org/psych\\_pract/treatg/pg/Practice%20Guidelines\\_8904/MajorDepressiveDisorder\\_2e.pdf](http://www.psych.org/psych_pract/treatg/pg/Practice%20Guidelines_8904/MajorDepressiveDisorder_2e.pdf). Accessed November 2, 2005.
27. Diagnostic and Statistical Manual of Mental Disorders, 4th ed. text rev. Washington, DC. *American Psychiatric Association*, 2000.
28. Practice guideline for the assessment and treatment of patients with suicidal behaviors. *Am J Psychiatry*. 2003;160 (Suppl. 11):1-60.
29. Intermountain Health Care. Depression 2006 Update. Available at: <https://kr.ihc.com/ext/Dcmnt?ncid=51061767>. Accessed September 19, 2007.
30. Brown GK, Bruce ML, Pearson JL. High-risk management guidelines for elderly suicidal patients in primary care settings. *Int J Geriatr Psychiatry*. 2001;16(6):593-601.
31. Lawlor DA, Hopker SW. The effectiveness of exercise as an intervention in the management of depression: systematic review and meta-regression analysis of randomized controlled trials. *Br Med J* 2001;322:1-8.
32. Dunn AL, Trivedi MH, O'Neal HA. Physical activity dose-response effects on outcomes of depression and anxiety. *Med Sci Sports Exercise* 2001;6 (Supp):S587-S597.
33. Blumenthal JA, Babyak MA, Doraiswamy PM, Watkins L, Hoffman BM, Barbour KA, Herman S, Craighead WE, Brosse AL, Waugh R, Hinderliter A, Sherwood A. Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosom Med*. 2007 Sep-Oct;69(7):587-96.
34. Blumenthal JA, Babyak MA, Moore KA, Craighead WE, Herman S, Khatri P, Waugh R, Napolitano MA, Forman LM, Appelbaum M, Doraiswamy PM, Krishnan KR. Effects of exercise training on older patients with major depression. *Arch Intern Med* 1999;159:2349-2356.
35. Duloxetine (Cymbalta): A new SNRI for depression. *Med Lett Drugs Ther*. 2004;46(1193):81-82.
36. FDA proposed antidepressant warning. Food and Drug Administration Web site. Available at: [www.fda.gov/cder/drug/antidepressants/default.htm](http://www.fda.gov/cder/drug/antidepressants/default.htm). Accessed July 15, 2007.
37. University of Michigan Health System. Depression: guidelines for clinical care. October 2005. Available at: <http://cme.med.umich.edu/pdf/guideline/depression04.pdf>. Accessed July 30, 2007.
38. Treatment Guidelines from the Medical Letter. Drugs for psychiatric disorders. *Treat Guidel Med Lett*. 2003;1(11):69-76.