



City Health Information

May 2008

The New York City Department of Health and Mental Hygiene

Vol. 27(5):33-40

RABIES IN NEW YORK CITY AND HUMAN RABIES PREVENTION

- When a patient presents with an animal bite, assess the need for rabies postexposure prophylaxis (PEP), and ensure that PEP is given properly when needed.
- Be aware that PEP is indicated only in particular circumstances, depending on whether a true exposure has occurred, the type of animal, and its availability for observation or testing. Raccoons and bats are the primary vectors of rabies in NYC.
- Report every animal bite to the NYC Health Department.

The rabies virus can infect all warm-blooded mammals, leading to acute, rapidly progressive encephalitis that results in death. When an animal is infected, the rabies virus moves along the peripheral nerves to the central nervous system (CNS) and brain, and then to the salivary glands. Initial signs appear after an incubation period of several weeks to months. These include a change in the animal's behavior; some animals may become aggressive and others may become reserved and withdrawn. Soon after, nonspecific symptoms such as lethargy, fever, and anorexia appear, followed by neurologic signs such as changes in gait, difficulty

swallowing, and paralysis and death within days. In almost all cases of human rabies, the virus is transmitted through the bite of a rabid animal.¹


Fortunately, in the United States (US), rabies in humans is extremely uncommon, and postexposure prophylaxis (PEP) is not automatically indicated when a person is bitten by an animal. It is important for the clinician to consider up-to-date epidemiologic data on rabies in NYC (**Resources**) when treating patients for bite wounds. From 2003 to 2006, there were 24,343 Emergency Department (ED) visits for animal bites reported in the 5 boroughs, largely due to dogs (73%), cats (13%), and rodents (7%). During that time, 2,464 animals were tested for rabies and 92 tested positive. Most (79%) of those were raccoons, predominantly from the Bronx and Staten Island. None were dogs.

EPIDEMIOLOGY OF RABIES

Animals

Different variants of the rabies virus exist and are maintained in their respective vectors, following geographically distinct patterns. In the US, those vectors include raccoons, skunks, bats, and foxes. The raccoon variant is present along the Atlantic Coast and the bat variant throughout the US except Hawaii. Most rabid animals in the US are raccoons, skunks, bats, and foxes.²



FIGURE 1. LABORATORY-CONFIRMED RABID ANIMALS IN NYC, 2007**TABLE 1. LABORATORY-CONFIRMED RABID ANIMALS IN NYC, 1992-2007**

Animal Species	Bronx	Queens	Manhattan	Staten Island	Brooklyn	Total Positive	Total Tested
Raccoon	144	7	8	117	0	276	5007
Skunk	33	0	0	2	0	35	197
Bat	7	4	7	4	4	26	712
Dog	0	0	0	0	0	0	1639
Cat	1	0	1*	7	1	10	2383
Other†	1	0	0	3	0	4	1714
Total	186	11	16	133	5	351	11652

*A cat from Columbia County that was tested in NYC.

†Other includes 1 coyote (1999, Bronx), 2 opossums (1994 and 1992, Staten Island), 1 groundhog (2007, Staten Island).

In NYC, rabies most often occurs in raccoons (**Figure 1, Table 1**). Occasionally, rabid raccoons transmit the virus to other animals, usually skunks. In NYC, rabid raccoons were first reported in 1992 in the Bronx and Staten Island, and are still largely limited to those areas, with isolated reports of rabid raccoons in Queens. A much smaller

number of rabid bats have been found in all 5 boroughs. Dogs and cats are also susceptible to rabies, but vaccination and animal control have significantly reduced the numbers of rabid dogs and cats. No rabid dogs have been reported in NYC since 1954. Since 1992, only 10 rabid cats have been identified in NYC; of these, 7 were

strays found in Staten Island in 2006 and 2007. Long Island was free of raccoon rabies until 2004, when the virus was identified in Nassau County; it has since spread to Suffolk County.

Rodents and members of the rabbit family are not considered to be vectors of rabies. The one exception is the groundhog, which has tested positive for rabies in NYC. Rabies infection of other domestic rodents or rabbits should be considered in unusual circumstances, such as when the animal has a bite wound and has been housed in an outdoor hutch, or when a rabies vector species (e.g., raccoon) has been observed in the vicinity.

Humans

In the US, an average of 1 to 2 cases of human rabies are reported each year, primarily due to exposure to rabid bats.³ Rabies has also been reported in travelers who become infected while overseas. Rarely, organ transplantation has resulted in transmission of rabies.^{4,5} Providers should ask about animal exposure and travel history for any patient with rapidly progressive encephalopathy.

There has not been a reported case of human rabies in NYC since the 1940s. Since 1992, 9 NYC residents have been bitten by animals known to be rabid (5 raccoons, 3 cats, 1 bat); all received rabies PEP and none developed rabies.

In the past 50 years, there have been 3 human cases of rabies in New York State; these occurred in a laboratory worker exposed while working with the virus,⁶ a traveler bitten by a rabid puppy while in Ghana,⁷ and an 11-year-old girl infected with the bat variant.⁸

Clinical Picture

The incubation period for human rabies is 31-90 days, although there have been rare reports of incubations of a year or more. Factors such as the extent of innervation of the bite site and its proximity to the CNS, severity of the bite, amount of inoculum, and immune status of the victim may affect incubation time.⁹

The first symptoms are often nonspecific (malaise, chills, fever, fatigue, respiratory problems, gastrointestinal distress) and may be attributed to common viral or bacterial infections. As the illness progresses, neurologic or psychiatric symptoms emerge. The one specific symptom reported by many patients is pain or paresthesias starting at the site of the bite and radiating out along the extremity or area of the healed bite wound. Eventually, CNS complications dominate the clinical picture. Findings may

include fever, paresthesias, stiff neck, muscle twitching, convulsions, hyperventilation, hydrophobia, and hypersalivation. The patient may exhibit hyperactivity, anxiety, agitation, hallucinations, thrashing, running, biting, or other bizarre behavior, as well as paralysis (20% of cases), and will ultimately become comatose and die.^{9,*}

Diagnostic Testing

Rabies in humans can be diagnosed by fluorescent antibody test on biopsy (neck skin) or corneal impressions, by virus isolation from saliva, virus neutralization assay on serum or cerebrospinal fluid (for evidence of rabies antibody), or reverse transcription polymerase chain reaction (RT-PCR) for viral RNA and genomic nucleotide sequence analysis on saliva. Human rabies testing is available at the Wadsworth Rabies Laboratory in Slingerlands, NY; special arrangements for transportation of critical specimens can be made through the NYC Health Department.¹⁰ The testing process requires extensive resources and should only be pursued when rabies is highly suspected based on clinical presentation, progression of illness, and history of animal exposure. All suspected cases of human rabies should be reported immediately, and the diagnosis should be pursued in consultation with the NYC Health Department Bureau of Communicable Disease (BCD).

DETERMINING EXPOSURE

Exposure to rabies is characterized as either a bite or non-bite exposure; each carries a different degree of risk.¹¹

- **Bite (higher-risk)** – Any penetration of skin by an animal's teeth. Bites to the face and hand and multiple bites carry the highest risk of rabies infection.
- **Non-bite (lower-risk)** – Scratches or abrasions received from an animal, or the contamination of open cuts, mucous membranes, or wounds with an animal's saliva (or brain and other neural tissue). Non-bite transmission of rabies is extremely rare.

Non-exposure: Petting or handling an animal; contact with blood, urine, or feces; and contact with saliva on intact skin do *not* constitute rabies exposures.

Possible exposure to a bat: Most bats found in NYC are not infected with rabies, but bats cause most cases of human rabies in the US. A bat bite causes minimal trauma, so that victims may not even be aware that a bite occurred. Because people have developed rabies after unrecognized

*One case of survival after intensive treatment of rabies infection has been documented (Willoughby RE, et al. *N Engl J Med.* 2005;35(24):2508-2514).

ASSESSING POSSIBLE EXPOSURE TO A BAT	
Exposure Unlikely	Exposure Reasonable Possibility
<ul style="list-style-type: none"> Bat droppings (guano) found in sleeping quarters. 	<ul style="list-style-type: none"> Bat found in same room with someone who might be unaware that exposure has occurred (e.g., sleeping, mentally disabled, or intoxicated person; unattended infant or child).
<ul style="list-style-type: none"> Teenager or adult touches a bat, but is certain the bat did not bite or scratch. 	<ul style="list-style-type: none"> Young child touches a live bat (may be unaware or unable to communicate exposure).
<ul style="list-style-type: none"> Bat swoops by a teenager or adult who does not feel it touch, or contact is not with bare skin (e.g., touches area covered by clothing). 	<ul style="list-style-type: none"> Bat flies into someone, touching bare skin and person is unable to determine what type of contact occurred.
<ul style="list-style-type: none"> Person has contact with a completely dried-up carcass of a bat. 	<ul style="list-style-type: none"> Someone with bare feet steps on a live bat.
<ul style="list-style-type: none"> Bats are heard or seen in walls or attic. 	<ul style="list-style-type: none"> Person puts hand in firewood or brush, feels pain, then sees a live bat.

bat exposures, reasonable possibility of exposure is treated as an exposure (see box to left).¹¹

HOW TO MANAGE POSSIBLE EXPOSURE TO RABIES

The need for rabies PEP is determined by assessing the exposure, the particular circumstances surrounding the bite, the type of animal, and its availability for rabies evaluation (**Table 2**). If the history suggests that an exposure has occurred and the animal cannot be observed or tested, PEP may be indicated (**Figure 2**). BCD is available for consultation (business hours: 212-788-9830; other times: 212-POISONS, 212-764-7667).

WHEN AND HOW TO ADMINISTER PEP

PEP consists of both rabies vaccine and human rabies immune globulin (HRIG), with certain exceptions (**Table 3**).

Rabies Vaccines


Two inactivated rabies vaccines are available in the US: Imovax[®], a human diploid cell vaccine (HDCV), and RabAvert[®], a purified chick embryo cell vaccine (PCEC). HDVC and PCEC are given in a 1-mL dose for both pre- and postexposure vaccination. A detectable antibody response occurs in 7-10 days, and immunity lasts at least 2 years. For adults, the vaccination should always be administered intramuscularly (IM) in the deltoid area. For children, the anterolateral aspect of the thigh is also acceptable. The gluteal area should never be used because administration in this area results in lower neutralizing antibody titers. The two licensed rabies

TABLE 2. MANAGING POTENTIAL RABIES EXPOSURES

- **Wound Care** – Evaluate and thoroughly irrigate and clean the wound. Assess need for tetanus immunization.
- **Exposure** – Determine whether there was a potential rabies exposure (see page 35).
- **History** – Get a good history including animal species, circumstances (provoked or not), owned vs. stray, location of animal, was animal ill or acting abnormally, vaccination status of animal.
- **Animal** – What type of animal? If an animal is capable of transmitting rabies, determine whether the animal is available for observation or testing.
- **Rabies PEP** – Give PEP if indicated (**Figure 2**).
- **Consult** – If needed, call BCD for consultation: 212-788-9830; non-business hours call 212-POISONS (212-764-7667).
- **Report** – Report the bite to the VPHS Animal Bite Unit at 212-676-2483.

BCD=Bureau of Communicable Disease; VPHS=Veterinary Public Health Services.

FIGURE 2. GUIDE TO RABIES POSTEXPOSURE PROPHYLAXIS



Was person EXPOSED*
to a dog, cat, raccoon, bat, skunk,
ferret, coyote, fox, groundhog,
livestock,¹ or other mammal?²
*Refer to definition of exposure,
page 35.

CONTACT THE NYC HEALTH DEPARTMENT
During regular business hours:
Veterinary Public Health Services to report all animal bites
at 212-676-2483.
The Bureau of Communicable Disease (BCD) for
consultation regarding rabies PEP at 212-788-9830.
During non-business hours call the Poison Control Center
at 212-764-7667.

Rodents and rabbits are not
considered to be vectors of
rabies. The one exception
is the groundhog, a rodent
that has tested positive for
rabies in NYC.

YES

NO
Rabies PEP
not indicated

Owned or Stray
Healthy Dog/
Cat/Ferret/Livestock

Raccoon, Skunk, Bat, Coyote, Fox, Groundhog
**Call BCD Immediately for
consultation: 212-788-9830**

Ill or Rabid-Acting Dog/Cat/
Livestock/Rabies Vector Species
**Call BCD Immediately for
consultation: 212-788-9830**

NO

**ANIMAL AVAILABLE
FOR 10-DAY
OBSERVATION?**

**ANIMAL AVAILABLE
FOR RABIES
TESTING?**

NO

**Consider Need
for Rabies PEP
ON A CASE-BY-CASE BASIS**
1) Additional factors
to consider include
circumstances of
exposure (provoked
or not), local
incidence of rabies,
health of animal.
2) If owned animal was
healthy and is familiar
to patient or patient's
guardian, consider
tracking down owner
to inquire about health
of animal before
starting PEP.
3) If animal was a healthy
stray, but is regularly
observed and readily
identifiable, consider
whether patient or
guardian may be able
to observe animal
before starting PEP.

YES

YES

10-DAY
OBSERVATION

RABIES TEST

Animal tests
POSITIVE

**RABIES
PEP
INDICATED**

Animal tests
NEGATIVE

Animal shows signs of
rabies or dies during
10-day observation

Consider *immediate*
rabies PEP if incident
is a high-risk head
wound, a severe
attack to a small child,
or if animal is highly
suspect for being
rabid even if animal
available for testing

**REMAINS
HEALTHY**

RABIES PEP NOT INDICATED

IMPORTANT POINTS TO REMEMBER
If animal is available or already undergoing observation or testing, counsel victim or guardian of victim regarding risk and need for PEP.
NO NEED TO INITIATE RABIES PEP IN THE FOLLOWING CIRCUMSTANCES:
1) Healthy dog, cat, ferret, or livestock undergoing 10-day observation.
2) Rabies vector species that is available for testing and results available within 24-48 hours.
3) Any animal considered to be rabid that is available for testing and results available within 24-48 hours.
Test results are available within 24 hours of submission to the Public Health Rabies Laboratory during regular business days. In some instances, emergency testing may be indicated and can be done in consultation with the NYC Health Department.

¹ Horses, mules, donkeys, sheep, goats, and swine (pigs).
² For exposures to other mammals, consult BCD.

vaccines are considered equally safe and effective. Reports in the literature indicate that interchanging vaccines is acceptable.^{11,12}

Serious problems from rabies vaccine are very rare. Mild problems include soreness, redness, swelling, or itching at the vaccination site (30%-74%); headache, nausea, abdominal pain, muscle aches, or dizziness (5%-40%). Moderate problems include hives, pain in the joints, fever (about 6% of booster doses), and (very rarely) illness resembling Guillain-Barré Syndrome with complete recovery.¹¹

Human Rabies Immune Globulin (HRIG)

HRIG must be administered to a patient receiving PEP unless the patient has had a complete rabies pre- or postexposure prophylaxis series in the past. HRIG neutralizes the virus at the wound site and reduces the amount of viable virus, which prevents or limits replication in muscle cells and subsequent invasion of nerve endings. Two versions of HRIG are available in the US: Imogram[®] and HyperRab[™]. HRIG dosing is weight-dependent. When possible, administer the full amount of HRIG in and around the wound site. For small children or limited injection sites (i.e., fingertip), inject as much HRIG around the site as possible; the remainder can be given IM at a site distant from vaccine administration (e.g., opposite deltoid muscle). Avoid administering in the gluteals if the patient is overweight.¹¹

Who Should Get Preexposure Vaccination

Preexposure vaccination is appropriate for people at high risk for rabies exposure, such as veterinarians, animal handlers, animal control workers, laboratory workers, and people planning travel to areas where rabies is endemic. Vaccine is given in three IM doses in the deltoid area on days 0, 7, and either day 21 or day 28. A booster should be provided according to ACIP guidelines (**Resources**). Preexposure vaccination does not eliminate the need for PEP, but it simplifies therapy (**Table 3**) and may offer protection to persons with unknown exposure or when PEP may be delayed.¹¹

Who Should Get Postexposure Vaccination

Only patients with a known or potential exposure to rabies should receive postexposure vaccination. The administration protocol depends on whether the patient has had prior immunization (**Table 3**). People who have received the vaccination series either pre- or postexposure are considered immunized and should adhere to the protocol for previously vaccinated patients. Rabies vaccination is costly and can be a traumatic experience for the recipient; therefore, it should not be given unless clearly indicated. If there is any doubt about the need for PEP or its proper administration, the clinician should consult BCD. Examples of issues that may arise in administering PEP are given in **Table 4**.^{11,12}

TABLE 3. RABIES POSTEXPOSURE PROPHYLAXIS REGIMENS

Regimen	Not Previously Vaccinated	Previously Vaccinated (either pre- or postexposure)*
Wound Cleansing	YES <ul style="list-style-type: none"> • Irrigation • Povidone-iodine • Determine need for tetanus 	YES <ul style="list-style-type: none"> • Irrigation • Povidone-iodine • Determine need for tetanus
Human Rabies Immune Globulin	YES <ul style="list-style-type: none"> • 20 IU/kg 	NO
Rabies Vaccine	YES <ul style="list-style-type: none"> • 1.0 mL IM in deltoid area[†] Days 0, 3, 7, 14, 28 	YES <ul style="list-style-type: none"> • 1.0 mL IM in deltoid area[†] Days 0 and 3

*Only applies to patients previously vaccinated using cell culture vaccine or any patient with a documented rabies virus neutralizing antibody titer.

†In children, anterolateral thigh can also be used. Avoid administration in gluteals.

More information on specific regimens is available at www.nyc.gov/html/doh/downloads/pdf/cd/acipmmwr-2008.pdf. IM=intramuscular.

TABLE 4. POSTEXPOSURE PROPHYLAXIS: WHAT TO DO IF...

- *A patient is off-schedule:* The PEP treatment schedule should be evaluated on a case-by-case basis.¹¹
 - DO NOT restart the series unless the patient began the series outside the US and the biologics used are questionable or no information is available. Consult the Bureau of Communicable Disease (BCD) for guidance.
 - HRIG should be given on day 0. Given after day 7, HRIG can interfere with the body's antibody production and may be contraindicated.
 - If the provider fails to administer HRIG on day 0, HRIG can be given up to 7 days after starting the vaccine schedule. If after day 7, contact BCD.
 - If HRIG was administered but NOT at the bite site, contact BCD.
 - Rabies vaccinations should be administered on days 0, 3, 7, 14, and 28. Deviations of a few days are of no great concern and the patient should resume the series, maintaining the recommended spacing between doses.
 - For example, if a patient misses day 3 and presents on day 5, the day 3 dose should be given that day, the day 7 dose on day 9, and the remaining doses on days 16 and 30.
 - If there is significant deviation from the schedule, resume series as above. Antibody titers should be assessed by RFFIT on a serum sample collected 14-28 days after finishing PEP to ensure a 1:5 serum dilution.
- *The patient is immunosuppressed at time of PEP:* Antibody titers should be assessed by RFFIT on a serum sample collected 14-28 days after finishing PEP to ensure a 1:5 serum dilution.¹¹
- *The patient is pregnant:* Pregnancy is not considered a contraindication to PEP; no fetal abnormalities have been associated with rabies vaccination.¹¹
- *The patient has begun PEP with a different rabies vaccine:* There are reports in the literature that indicate HDCV and PCEC vaccines can be interchanged.¹² If the original vaccine type is not available, continue vaccination with vaccine that is available.

RFFIT = rapid fluorescent focus inhibition test.

Most EDs in NYC stock rabies biologics and typically serve as the appropriate health care site for patients to initiate the PEP series. Every effort should be made to ensure that patients receive all doses at the appropriate intervals.

Physicians can order biologics directly from the manufacturer. The Health Department does not provide rabies biologics or administer PEP directly.

Animal Bite Follow-Up

The NYC Health Code requires that all animal bites be reported. The NYC Office of Veterinary Public Health Services (VPHS) receives those reports and coordinates follow-up with the animal, its owner, and the person bitten to ensure that the animal is managed appropriately and the public is protected from further harm. VPHS arranges animal capture and animal rabies testing if indicated. To report an animal bite or provide information about the biting animal, call the VPHS Animal Bite Unit at 212-676-2483, from 9 a.m. to 5 p.m., Monday to Friday, or the Poison Control Center at 212-POISONS (212-764-7667) all other times.

INFORMATION FOR VETERINARIANS

NYC Health Code Section 11.64 requires that animal diseases that are communicable to humans and that are of public health concern be reported to BCD (see box on back). Influenza, which was recently added to the list, is specified as any strain of influenza with the potential to cause a pandemic. Influenza viruses can be found in several types of animals including birds, pigs, equines, canines, and sea mammals.

To make a report during regular business hours, call BCD at 212-788-9830. For urgent matters outside regular business hours, call the Poison Control Center at 212-POISONS (212-764-7667). For more information about animal disease reporting, visit the Health Department Web site at: www.nyc.gov/html/doh/html/zoo/zoo-reporting.shtml.

SUMMARY

When treating patients with reported animal bites or other potential rabies exposures, advise them that the risk of rabies transmission is low except for certain situations (animal bites from raccoons, bats, skunks, and other rabies vector

species). For dog or cat bites, rabies postexposure prophylaxis is usually not indicated. Whenever possible, the animal should be observed and/or tested. When rabies PEP is indicated, it is essential that patients are followed closely to ensure that they complete preventive therapy. ♦

REPORTABLE ANIMAL DISEASES

Report if disease is suspected

- Rabies
- Anthrax
- Brucellosis
- Glanders
- Plague
- Q fever
- Tularemia
- Monkeypox
- Influenza caused by novel viral strain with pandemic potential
- Severe acute respiratory syndrome (SARS)
- Arboviral encephalitis, acute

Report if disease is diagnosed

- Psittacosis
- Leptospirosis

RESOURCES

- NYC Health Department Rabies Web site www.nyc.gov/html/doh/html/cd/cdrab.shtml
- New York City Health Department Veterinary Public Health Services www.nyc.gov/html/doh/html/vet/vet1.shtml
- ACIP Guidelines on the Prevention of Rabies in Humans www.nyc.gov/html/doh/downloads/pdf/cd/acip-mmwr-2008.pdf
- NASPHV Animal Rabies Compendium www.nyc.gov/html/doh/downloads/pdf/zoo/zoo-rabies-compendium-2008.pdf
- CDC Rabies Web site www.cdc.gov/rabies

HOW TO CONTACT THE NYC HEALTH DEPARTMENT

- Bureau of Communicable Disease: 212-788-9830.
 - Veterinary Public Health Services: 212-676-2483.
- During non-business hours call the Poison Control Center at 212-764-7667.

Resource listings are provided for informational purposes only and do not imply endorsement by the NYC DOHMH.

References Available Online: www.nyc.gov/html/doh/downloads/pdf/chi/chi27-5.pdf.

RECEIVE CHI BY E-MAIL Each time **City Health Information** is published you will receive a link to the issue in PDF format. To subscribe, visit www.nyc.gov/html/doh/html/chi/chi.shtml.

DOHMH JOB OPENINGS: We seek doctors, nurses, administrators, social workers, and other public health professionals. Visit www.nyc.gov/health/careers to view openings.



City Health Information



May 2008 The New York City Department of Health and Mental Hygiene Vol. 27(5):33-40

2 Lafayette Street, 20th Floor, CN-65, New York, NY 10007 (212) 676-2188

Michael R. Bloomberg
Mayor

Thomas R. Frieden, MD, MPH
Commissioner of Health and Mental Hygiene

Division of Epidemiology
Lorna E. Thorpe, PhD, Deputy Commissioner

Division of Disease Control
Isaac Weisfuse, MD, MPH, Deputy Commissioner

Bureau of Communicable Disease
Marcelle Layton, MD, Assistant Commissioner

Zoonotic, Influenza, and Vectorborne Disease Unit
Annie D. Fine, MD, Director
Sally A. Slavinski, DVM, MPH, Assistant Director

Bureau of Public Health Training
Carolyn Greene, MD, Assistant Commissioner
Peggy Millstone, Director, Scientific Communications
Rita Baron-Faust, CHES, Medical Editor
Peter Ephross, Medical Editor
Quawana Charlton, Editorial Assistant

Copyright ©2008 The New York City Department of Health and Mental Hygiene
Email *City Health Information* at: nycdohrp@health.nyc.gov
Suggested citation: Slavinski SA, Fine AD. Rabies in New York City and human rabies prevention. *City Health Information*. 2008;27(5):33-40.

PRST STD
U.S. POSTAGE
PAID
NEW YORK, N.Y.
PERMIT NO. 6174

RABIES REFERENCES

REFERENCES

1. Centers for Disease Control and Prevention. Compendium of animal rabies prevention and control, 2007. *MMWR*. 2007;56(RR03):1-8.
2. Blanton JD, Hanlon CA, Rupprecht CE. Rabies surveillance in the United States during 2006. *J Am Vet Med Assoc*. 2007;231(4):540-556.
3. Centers for Disease Control and Prevention. Rabies. <http://cdc.gov/rabies>. Accessed April 13, 2008.
4. Centers for Disease Control and Prevention. Investigation of Rabies Infections in Organ Donor and Transplant Recipients—Alabama, Arkansas, Oklahoma, and Texas, 2004. *MMWR*. 2004;53(26):586-589.
5. Srinivasan A, Burton EC, Kuehnert MJ. Transmission of rabies virus from an organ donor to four transplant recipients. *N Engl J Med*. 2005;352(11):1103-1111.
6. Centers for Disease Control and Prevention. Rabies in a laboratory worker, New York. *MMWR*. 1977;26:183-184.
7. Centers for Disease Control and Prevention. Human rabies—New York, 1993. *MMWR*. 1993;42(41):799,805-806.
8. Centers for Disease Control and Prevention. Human rabies—California, Georgia, Minnesota, New York, and Wisconsin, 2000. *MMWR*. 2000;49:1111-1115.
9. Baer G, ed. *The Natural History of Rabies*. 2nd ed. Boca Raton, FL: CRC Press Inc; 1991.
10. Wadsworth Center. Antemortem diagnosis of human rabies. www.wadsworth.org/rabies/prof/ante.htm. Accessed April 14, 2008.
11. Centers for Disease Control and Prevention. Human rabies prevention—United States, 2008. Recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR*. 2008;57(RR3):1-28. www.nyc.gov/html/doh/downloads/pdf/cd/acip-mmwr-2008.pdf. Accessed May 13, 2008.
12. Briggs DJ, Dreesen DW, Nicolay U, et al. Purified chick embryo cell culture rabies vaccine: interchangeability with human diploid cell culture rabies vaccine and comparison of one versus two-dose post-exposure booster regimen for previously immunized persons. *Vaccine*. 2000(9-10);19:1055-1060.

Continuing Education Activity

Rabies in New York City and Human Rabies Prevention

SPONSORED BY

THE NEW YORK CITY DEPARTMENT OF
HEALTH AND MENTAL HYGIENE (DOHMH)
CITY HEALTH INFORMATION
MAY 2008 VOL 27(5):33-40

Objectives

At the conclusion of this activity, participants should:

1. Be familiar with the epidemiology of rabies in NYC and risks for exposure to NYC residents.
2. Understand when rabies postexposure prophylaxis (PEP) is indicated.
3. Know proper administration of PEP which includes Human Rabies Immune Globulin (HRIG) and 5 doses of rabies vaccine according to schedule.
4. Understand the proper management of animals that have bitten and the role of the DOHMH.

Accreditation

New York City Department of Health and Mental Hygiene (NYC DOHMH) is accredited by the Medical Society of the State of New York to sponsor continuing medical education for physicians. The NYC DOHMH designates this continuing medical education activity for a maximum of 2.00 AMA PRA Category 1 credit(s)[™]. Each physician should only claim credit commensurate with the extent of his/her participation in the activity.

Participants are required to submit name, address, and professional degree. This information will be maintained in the Department's CME program database. If you request, the CME Program will verify your participation and whether you passed the exam.

We will *not* share information with other organizations without your permission, except in certain emergencies when communication with health care providers is deemed by the public health agencies to be essential or when required by law. Participants who provide e-mail addresses may receive electronic announcements from the Department about future continuing education activities as well as other public health information.

Participants must submit the accompanying exam by May 30, 2009.

CME Faculty:

Sally A. Slavinski, DVM, MPH
Annie D. Fine, MD

All faculty are affiliated with the NYC DOHMH. The faculty does not have any financial arrangements or affiliations with any commercial entities whose products, research, or services may be discussed in this issue.

CME Activity Rabies in New York City and Human Rabies Prevention

May 2008

1. How is rabies transmitted?

- A. A bite from a rabid animal.
- B. Touching the fur of a rabid animal.
- C. Being in the same room as a rabid animal.
- D. Handling urine of an infected animal.
- E. Exposing mucous membranes to infective saliva or neurologic tissue.
- F. A and E.

2. If a person is bitten by a dog that appears healthy but is not vaccinated against rabies, what should be done?

- A. Rabies postexposure prophylaxis should ALWAYS be initiated.
- B. Do not initiate rabies postexposure prophylaxis if the dog remains healthy and can be observed 10 days.
- C. Report the animal bite to the DOHMH.
- D. A and C.
- E. B and C.

3. What animal is responsible for the most human rabies cases in the United States every year?

- A. Raccoons.
- B. Dogs.
- C. Cats.
- D. Bats.

4. When administering rabies postexposure prophylaxis, under which circumstances is human rabies immune globulin (HRIG) NOT indicated?

- A. The patient has received a complete rabies preexposure or postexposure rabies vaccination series in the past.
- B. The patient doesn't want it.
- C. The animal bite occurred more than a week earlier.
- D. The exposure was from a bat.

5. For which of the following scenarios is rabies postexposure prophylaxis indicated?

- A. A 43-year-old woman is bitten by her best friend's healthy dog in Queens.
- B. A 10 year old child who woke to find bats flying in her tent while at summer camp in upstate NY.
- C. A 26-year-old man who was bitten by a raccoon that he trapped in the Bronx yesterday, which is now being tested for rabies at the DOHMH Public Health Laboratory.
- D. A 33-year-old woman who was bitten by a stray cat that attacked her dog while she was walking it in Staten Island. The cat cannot be found.
- E. A and B.
- F. B and D.

6. How well did this continuing education activity achieve its educational objectives?

- A. Very well
- B. Adequately
- C. Poorly

PLEASE PRINT LEGIBLY.

Name _____ Degree _____

Address _____


City _____ State _____ Zip _____

Date _____ Telephone _____

E-mail address _____



NO POSTAGE
NECESSARY
IF MAILED
IN THE
UNITED STATES



BUSINESS REPLY MAIL
FIRST-CLASS MAIL PERMIT NO.2379 NEW YORK NY

POSTAGE WILL BE PAID BY ADDRESSEE

CME ADMINISTRATOR
NYC DEPARTMENT OF HEALTH
AND MENTAL HYGIENE
2 LAFAYETTE ST, CN - 65
NEW YORK, NY 10277-1632



Continuing Education Activity

This issue of *City Health Information*, including the continuing education activity, can be downloaded at www.nyc.gov/html/doh/html/chi/chi.shtml.

Instructions

Read this issue of *City Health Information* for the correct answers to questions. To receive continuing education credit, you must answer 4 of the first 5 questions correctly.

To Submit by Mail

1. Complete all information on the response card, including your name, degree, mailing address, telephone number, and e-mail address. PLEASE PRINT LEGIBLY.
2. Select your answers to the questions and check the corresponding boxes on the response card.
3. Return the response card (or a photocopy) postmarked **no later than May 30, 2009**. Mail to:

CME Administrator, NYC Dept. of Health and Mental Hygiene,
2 Lafayette Street, CN-65, New York, NY 10277-1632.

To Submit Online

Visit www.nyc.gov/html/doh/html/chi/chi.shtml to submit a continuing education test online. Once logged into NYC MED, use the navigation menu in the left column to access this issue of *City Health Information*. Your responses will be graded immediately, and you can print out your certificate.