



PARENTS!



Protect your child from these serious diseases that can cause your child to become very ill or hospitalized. These diseases may cause blindness, deafness, brain damage or in rare cases, death...

- | | | |
|----------------------------|------------------------|-------------------------------|
| Hepatitis B | Measles | Hepatitis A |
| Polio | Mumps | Meningitis |
| Tetanus | Rubella | Rotavirus |
| Diphtheria | Varicella (Chickenpox) | Haemophilus Influenzae type b |
| Pertussis (Whooping Cough) | Pneumococcus | Influenza |

Vaccines...




- *Protect your child against these diseases*
 - *Help your child develop immunity without getting sick*
 - *Are safe and effective*
 - *Are required for your child to attend day care and school*

“Start your baby’s shots on time and stay on schedule”

See your child’s doctor, or for information on where your child can be vaccinated, Dial 311, or visit nyc.gov/health

RECOMMENDED PRESCHOOL IMMUNIZATION SCHEDULE, JANUARY 2010

	BIRTH	2 MONTHS	4 MONTHS	6 MONTHS	12 MONTHS	15 MONTHS	4-6 YEARS
Hepatitis B HepB	√	√		√ 6-18 months			
Diphtheria, Tetanus, Pertussis DTaP		√	√	√		√ 15-18 months	√
Haemophilus influenzae type b Hib		√	√	√	√ 12-15 months		
Polio IPV		√	√	√ 6-18 months			√
Pneumococcal Disease PCV		√	√	√	√ 12-15 months		
Rotavirus RV		√ 6-14 weeks	√	√ 8 months			
Measles, Mumps, Rubella MMR					√		√
Varicella Var					√		√
Hepatitis A HepA					2 doses 12-23 months	The 2 nd dose should be given 6 months after the 1 st dose	
Influenza				Yearly - Children under 9 should receive 2 doses in the first year of vaccination			
H1N1 Influenza				2 doses for ages 9 and younger. 2 nd dose should be given 3 weeks after the 1 st dose			
Meningococcal MCV4						Children 2 years and older in certain high-risk groups	



IMM 43 (05/10)



Health

Bureau of Immunization

