

Fentanyl causes overdose deaths.

**Anyone using heroin, cocaine,
crack or pills from nonmedical sources,
even occasionally, is at risk.**

Reduce your overdose risk:

- Test your drugs using fentanyl test strips.
 - If positive, consider not using. If you do plan to use, start with a small amount and go slowly.
- Avoid using alone. Take turns or call the Never Use Alone hotline at 800-484-3731.
- Avoid mixing drugs.
- Keep naloxone ready and on hand.

For mental health support or resources, call **888-692-9355**.

For more information, visit **nyc.gov/overdose**.

Fentanyl lakòz moun mouri nan ovèdoz.

Nenpòt moun k ap itilize ewoyin,
kokayin, krak oswa grenn ki pa medikal,
menm si se yon fwa konsa, gen risk.

Diminye risk pou w fè ovèdoz:

- Teste dwòg ou yo gras ak bann tèz Fentanyl yo.
 - Si tèz yo ta pozitif, pa itilize dwòg lan. Si ou planifye pou itilize l, kòmanse ak yon ti kantite epi ale lantman.
- Evite itilize l poukont ou. Pa fè sa yon sèl kou oswa ou kapab rele liy asistans Never Use Alone (Pa Janm Itilize l Poukont Ou) lan nan 800-484-3731.
- Evite melanje dwòg.
- Toujou mete naxolone yon kote tou pare ou ka pase men w pran l.

Pou sipò oswa resous ki gen pou wè ak sante mantal, rele nan **888-692-9355**.

Pou jwenn plis enfòmasyon, ale sou **nyc.gov/overdose**.