

**Fentanyl causes overdose deaths.**

**Anyone using heroin, cocaine,  
crack or pills from nonmedical sources,  
even occasionally, is at risk.**

Reduce your overdose risk:

- Test your drugs using fentanyl test strips.
  - If positive, consider not using. If you do plan to use, start with a small amount and go slowly.
- Avoid using alone. Take turns or call the Never Use Alone hotline at 800-484-3731.
- Avoid mixing drugs.
- Keep naloxone ready and on hand.

For mental health support or resources, call **888-NYC-WELL** (888-692-9355).

For more information, visit **[nyc.gov/overdose](https://nyc.gov/overdose)**.

**펜타닐은 과다 복용으로 인한 사망을 유발합니다.**

**헤로인, 코카인, 크랙 또는  
비의료용 알약을 사용하는 사람은 누구나  
때때로 위험에 처할 수 있습니다.**

과다복용 위험을 줄이는 방법:

- 펜타닐 테스트 스트립을 사용하여 약물을 검사하십시오.
  - 양성이라면 사용하지 않는 것이 좋습니다. 사용할 계획이 있다면 소량으로 천천히 시작하십시오.
- 단독으로 사용하지 마십시오. 교대로 사용하거나 단독 사용 금지(Never Use Alone) 핫라인에 800-484-3731 번으로 전화하십시오.
- 약물을 섞지 마십시오.
- 날록손을 준비해 두십시오.

정신 건강 지원 또는 리소스는 **888-NYC-WELL (888-692-9355)**  
번으로 문의하십시오. 자세한 내용은 [nyc.gov/overdose](https://nyc.gov/overdose)를 참조하십시오.