

**PRESCRIBE NALOXONE TO ANYONE WHO REQUESTS IT, INCLUDING:**

- ✓ Anyone at risk of experiencing an opioid-related overdose
- ✓ Anyone, including friends and family, who may assist someone at risk for an opioid-related overdose
- ✓ Anyone receiving prescription opioid analgesics

**Providers may wish to conduct a risk assessment to identify *additional* patients who need naloxone.**

**The following factors increase overdose risk and may serve as additional screening criteria:**

- High-dose opioid prescription (90 or more daily morphine milligram equivalents); chronic opioid therapy (for three months or longer); and concurrent opioid and benzodiazepine prescriptions
- Current or previous opioid misuse or illicit drug use; treatment for opioid use disorder (e.g., methadone, buprenorphine, naltrexone); and opioid overdose history

## Narcan® Nasal Spray

4 mg (NDC 69547-353-02)  
1 x two-pack  
Refills: PRN


Sig: For suspected opioid overdose  
Follow package instructions.



## Intramuscular

Naloxone HCl 0.4 mg/mL  
2 x 1 mL single-dose vials  
(NDC 0409-1215-01 or 67457-292-02)  
Refills: PRN  
2 x 23 g, 3 mL,  
1-inch syringe  
Refills: PRN

Sig: For suspected opioid overdose  
Inject 1 mL intramuscularly in shoulder or thigh.




## Intranasal

Naloxone HCl 1 mg/mL  
2 x 2 mL as pre-filled Luer-Lock syringe  
(NDC 76329-3369-1)  
Refills: PRN  
2 x intranasal mucosal atomizing device  
(MAD 300)  
Refills: PRN

Sig: For suspected opioid overdose  
Spray 1 mL in each nostril.  
If no response in two minutes, give second dose.

Atomizer typically not covered by insurance.

*Note to pharmacist:* Call 866-246-6990 or  
800-723-3892 to order MAD 300.



## Evzio® Auto Injector

0.4 mg (NDC 60842-030-01)  
1 x two-pack  
Sig: For suspected opioid overdose  
Follow package instructions.

Prior authorization is typically required.

