

COVID-19 Infection Prevention Competency Tools for Use in Congregate Residential Facilities

Introduction

Congregate residential settings present unique infection control challenges. A congregate residential setting is a facility where a group of people reside and gather in close proximity with each other for a limited or extended period of time. This includes nursing homes, adult care facilities, group homes, homeless shelters, correctional facilities, residential substance use treatment facilities, and supportive housing sites. Interventions to prevent and interrupt COVID-19 transmission should be part of a comprehensive infection surveillance plan for your facility. The competency tools included in this document focus on cleaning and disinfecting as well as the use of personal protective equipment (PPE), and serve as a guide to evaluate infection prevention processes at your facility.

How to use the competency tools

A staff member may act as the evaluator and observe another staff member perform the sequence detailed in the tables below on an annual basis, or more frequently if deviations from practice are identified at your facility. The evaluator should be a senior or manager level staff member and have knowledge of the processes in the competencies and can teach the process being evaluated. The evaluator guidance in the competencies below serves as a prompt to the evaluator to look for specific details or to ask questions of the staff being evaluated. The staff member being evaluated should be asked to verbalize each step as it is performed. If any step is not completed properly, the staff member should be instructed on how to properly complete the task. Depending on the overall outcome, the evaluation should be repeated at a later date at the discretion of the evaluator (i.e., a few weeks later) to assess improvement. When each step is completed properly, the staff member should be considered competent in that activity. For assistance using the competency tools or other infection prevention questions, contact the NYC Health Department's Congregate Settings Investigation and Response Unit infection prevention team at IPC@health.nyc.gov.

7.20.21

Personal Protective Equipment (PPE) Cleaning and Disinfecting Competency Tool¹

Employee Name:	Job Title:
Evaluator Name:	Job Title:

Task	Evaluator Guidance		etent
I d5K	Evaluator Guidance	Yes	No
Gather appropriate supplies	 Appropriate disinfectant(s) listed by the Environmental Protection Agency (EPA) as effective against COVID-19². Additional cleaning supplies as needed (e.g., cleaning rag). Appropriate PPE (e.g., gloves and eye protection). 		
Don (put on) PPE when appropriate	When cleaning a room occupied by a resident with confirmed or suspected COVID-19: • Don full PPE (gown, gloves, eye protection, and N95 or higher level respirator) in the correct sequence. For routine cleaning: • Don gloves. Note: Staff should not touch personal items such as their cell phone with soiled gloves and when in PPE.		
Clean soiled area	Use soap and water or other cleaning agent to remove visible dirt or buildup.		

¹ Centers for Disease Control and Prevention (2021, June 15) Cleaning and Disinfecting Your Facility

² Environmental Protection Agency (EPA) About List N: Disinfectants for Coronavirus (COVID-19)

³ The correct sequence for donning PPE is as follows: (1) identify and gather the proper PPE. (2) Perform hand hygiene. (3) Put on isolation gown. (4) Put on N95 or higher level respiratory (facemask if respirator not available). (5) Put on eye protection. (6) Put on gloves. Source: <u>Centers for Disease Control and Prevention (2020, June 9)</u>
<u>Using Personal Protective Equipment (PPE)</u>

	Note: Not applicable for electronics. For		
	electronics, check manufacturer's cleaning		
	recommendations.		
Disinfect area	Disinfect surface with appropriate disinfectant		
	following instructions on the label.		
	Ask participant to verbalize:		
	 Manufacturer's cleaning instruction for 		
	electronics and other objects		
	 The disinfectant's contact time (how long 	5	
	the surface must remain wet for the		
	disinfectant to be effective).		
Doff (take off) PPE	After cleaning a room used by a symptomatic		
	person:		
	 Refer to doffing sequence in competency 		
	tool.		
Dispose of trash	Properly dispose of any trash generated from		
	the cleaning and disinfection process.		
	A clean set of gloves is donned prior to		
	touching and moving the trash.		
Remove gloves safely	Hand hygiene can be performed with either		
and perform hand	hand sanitizer or soap and water. Observe		
hygiene	individual perform hand hygiene properly		
	including the front and back of hands, in		
	between the fingers, fingertips, thumb, and wrist area. ⁴		
	wrist area.		
Follow up actions:			
Tonow up detions:			
Employee Signature:	Date	e:	
Evaluator Signature:	Date	e:	

⁴ World Health Organization (2009, May) How to Handwash?

Personal Protective Equipment (PPE) Donning when Caring for a Resident with Confirmed or Suspected COVID-19 Competency Tool

Employee Name:	Job Title:
Evaluator Name:	Job Title:

Took	Fredricker Cridenes	Competent	
Task	Evaluator Guidance	Yes	No
Gather appropriate supplies	 Supplies include: 1 pair of gloves Disposable gown 1 N95 respirator (or equivalent or higher level) or disposable mask if respirator is not available Eye protection (face shield or goggles) 		
	Note: The individual should identify the type of respirator and size for which they have been previously <u>fit-tested or other appropriate mask</u> .		
Perform hand hygiene	Hand hygiene can be performed with either hand sanitizer or soap and water. Observe individual perform hand hygiene properly including the front and back of hands, in between the fingers, fingertips, thumb, and wrist area for 20 seconds.		
Don (put on) disposable gown	Opening of the gown goes in the back. Gown fits properly:		
Tie gown at the back and neck	Individual can ask for assistance to tie straps in the back. Once secured, the gown should cover the back and not slip from shoulders with movement.		
Don (put on) disposable mask or N- 95	 For surgical mask: Secure ties at crown of the head and base of the neck or hook loops around ears. Fit band to nose bridge. 		

	• Fit bottom of mask below the chin.		
	For N95:		
	Individual has the appropriate size and		
	mask.		
	Secure bottom band around base of neck.		
	Secure top band at crown of head. Do not		
	cross bands.		
	Fit band to nose bridge.		
	Perform fit check by:		
	 Exhaling into the respirator and 		
	feeling for air leaking around the		
	edges of the mask or		
	By inhaling so that the respirator All pages slightly had breath for 10.		
	collapses slightly, hold breath for 10		
	seconds, feeling for a slight collapse		
	of the respirator and air leakage around the edges of the mask		
Don eye protection	Face shield, goggles, or safety glasses are		
Don'eye protection	acceptable eye protection. Adjust fit as needed.		
Don gloves	Put on gloves and extend over the cuff of isolation	П	
Don gloves	gown.		
	80		
Follow up actions:			
	_		
Employee Signature:	Date:		
Limpioyee Signature	Date.		
Evaluator Signature:	Date:		

Additional resources:

- <u>Centers for Disease Control and Prevention (2020, August 19) Using Personal Protective</u> <u>Equipment (PPE)</u>
- <u>Centers for Disease Control and Prevention (2020, July 14) Demonstration of Donning</u> (Putting On) Personal Protective Equipment (PPE) video

Personal Protective Equipment (PPE)	
Doffing Competency Tool	

Employee Name:	Job Title:
Evaluator Name:	Job Title:

Note: Remember that the front of the gown, outside of gloves, and outside of eye protection are all considered contaminated. Any breaches in the PPE doffing process could result in self-contamination, and all steps should be performed with care.

Deffine Commons	Fundamenton Cuidones	Competent	
Doffing Sequence	Evaluator Guidance	Yes	No
Remove gown	Untie/unfasten at waist first, then at the neck.		
Remove gloves	 Grasp outside of one glove with opposite gloved hand and peel off. Hold removed glove in remaining gloved hand. Slide fingers of ungloved hand under remaining glove at the wrist. Peel off remaining glove and turn inside out, continuing to hold the first glove. Note: Gloves should be removed in a way that avoids contaminating the hands. The outside (contaminated side) of the glove should never touch the skin. There are multiple ways to remove gloves safely. This competency describes one way to remove gloves. If the individual fails to remove gloves safely, counsel them on technique. 		
Dispose gloves in trash			
Perform hand hygiene	Hand hygiene can be performed with either hand sanitizer or soap and water. Observe individual perform hand hygiene properly including the front and back of hands, in		

	between the fingers, fingertips, thumb, and	I	
	wrist area for 20 seconds.		
Remove eye protection/	Both items should be removed by the strap		
face shield	at the back of the head. Remove away from	ו	
	face.		
Remove mask or	 Do not touch the front of the 		
respirator	respirator or mask.		
	 For N95: remove the bottom strap 		
	by touching only the strap and bring	g	
	it carefully over the head. Grasp the	:	
	top strap and bring it carefully over		
	the head, and then pull the		
	respirator away from the face		
	without touching the front of the		
	respirator.		
	Note: Mask or respirator may be left on as		
	appropriate. If the mask or respirator has		
	become wet, damaged, or soiled, it should		
D ()	be removed and discarded.		
Perform hand hygiene	Hand hygiene can be performed with either	r 🗆	
	hand sanitizer or soap and water. Observe		
	individual perform hand hygiene properly		
	including the front and back of hands, in	.	
	between the fingers, fingertips, thumb, and wrist area for 20 seconds.	'	
	wrist area for 20 seconds.		
Follow up actions:			
Employee Signature		Datas	
Employee Signature:		Date:	
Evaluator Signature		Date:	

Additional resources:

• <u>Centers for Disease Control and Prevention (2020, August 19) Using Personal Protective</u> <u>Equipment (PPE)</u> • Centers for Disease Control and Prevention PPE doffing video (2020, April 21)
Demonstration of Doffing (Taking Off) Personal Protective Equipment (PPE) video

The NYC Health Department may change recommendations as the situation evolves. 7.20.21