

HIV Among People Born Outside the United States – New York City, 2024

HIV Epidemiology Program
New York City Department of Health and Mental Hygiene
Published December 2025
nyc.gov/hivreports



Table of Contents

Description	Slide
HIV – New York City	
by place of birth	
number of people newly diagnosed	5
number of people newly diagnosed and percent change	6
Basic statistics of HIV among people born outside the U.S. – New York City	7
HIV among people born outside the U.S. – New York City	
number of people newly diagnosed	8
number of people newly diagnosed and percent change	9
by gender	
number of people newly diagnosed	10
number of people newly diagnosed and percent change	11
by race and ethnicity	
number of people newly diagnosed	12
number of people newly diagnosed and percent change	13
by age group	
number of people newly diagnosed	14
number of people newly diagnosed and percent change	15
by race and ethnicity and age group	
number of people newly diagnosed	16
by borough	
number of people newly diagnosed	17
number of people newly diagnosed and percent change	18


Table of Contents

Description	Slide
HIV among people born outside the U.S. – New York City (continued)	
by neighborhood poverty level	
number of people newly diagnosed	19
number of people newly diagnosed and percent change	20
by transmission category	
number of people newly diagnosed	21
number of people newly diagnosed and percent change	22
Initiation of care within 30 days of diagnosis among people born outside the U.S. – New York City	
by demographic group	24
by United Hospital Fund Neighborhood	25
Viral suppression within three months of diagnosis among people born outside the U.S. – New York City	
by demographic group	26
by United Hospital Fund Neighborhood	27
Viral suppression among people born outside the U.S. and diagnosed with HIV – New York City	
by demographic group	28
by United Hospital Fund Neighborhood	29
Viral suppression among people born outside the U.S. and diagnosed with HIV – New York City	
by demographic group	31
by United Hospital Fund Neighborhood	32
	33

Table of Contents


Description	Slide
HIV care continuum among people born outside the U.S. – New York City	
overall and by race and ethnicity	34
Age-adjusted death rate per 1,000 people born outside the U.S. and diagnosed with HIV – New York City	
by demographic group	35
by United Hospital Fund Neighborhood	36
Deaths among people born outside the U.S. and diagnosed with HIV by cause of death – New York City	
	37
	38
Appendices	
How to find our data	39
Technical notes	40
Technical notes on the HIV care continuum	42
Reporting HIV and AIDS diagnoses for health care providers	43
Acknowledgements	44

Number of People Newly Diagnosed With HIV by Place of Birth – New York City, 2019-2024



Among people newly diagnosed with HIV, the difference in the number of people born in the United States and those born outside of the United States narrowed markedly from 2019 to 2024. The number of people newly diagnosed with HIV increased among people born outside of the United States and among people with an unknown place of birth from 2019 to 2024. The number of people newly diagnosed with HIV decreased for all other places of birth. In 2024, people born outside of the United States represented 35% of new HIV diagnoses overall and 47% among those with a known place of birth.

Number of People Newly Diagnosed With HIV and Percent Change by Place of Birth – New York City in 2019, 2023, and 2024



From 2019 to 2024, the number of people newly diagnosed with HIV increased among people born outside of the United States (23%) and among people with an unknown place of birth (111%); all other groups experienced decreases. From 2023 to 2024, an increase occurred among people with an unknown place of birth (32%). Due to the relatively large number of people with an unknown place of birth, percent change calculations by place of birth should be interpreted with caution.

Basic Statistics of HIV Among People Born Outside the U.S. – New York City, 2024

- **622 people newly diagnosed with HIV¹ who were born outside the United States**
 - Including 165 people concurrently diagnosed with AIDS² (26.5% of diagnoses)
- **428 people newly diagnosed with AIDS³ who were born outside the United States**
- **There are an estimated 23,800 people with HIV⁴ who were born outside the United States**
- **260 deaths⁵ among people with HIV who were born outside the United States**
 - 6.8 deaths per 1,000 people with HIV⁶

¹Excludes people known to have been diagnosed outside NYC.

²An AIDS diagnosis within 31 days of an HIV diagnosis is considered a concurrent diagnosis.

³Includes people concurrently diagnosed with HIV and AIDS.


⁴Approximate value calculated as the number of people with HIV divided by the estimated proportion of people with HIV who had been diagnosed, see Technical Notes for more details.

⁵Includes deaths from any cause in people with HIV. Death data for 2024 are incomplete.

⁶Age-adjusted to the 2000 U.S. Standard Population. People newly diagnosed with HIV at death were excluded from the analysis.


As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Number of New HIV Diagnoses Among People Born Outside the U.S. – New York City, 2019-2024




In 2024, people born outside of the U.S. represented 35% of new diagnoses overall in New York City. The number of people newly diagnosed with HIV who were born outside the U.S. decreased from 2019 to 2020 and increased from 2020 to 2024.

Number of New HIV Diagnoses and Percent Change Among People Born Outside the U.S. – New York City in 2019, 2023, and 2024




The number of people newly diagnosed with HIV who were born outside the U.S. increased by 23% from 2019 to 2024 in New York City. The number of people newly diagnosed from 2023 to 2024 was relatively flat.

Number of New HIV Diagnoses Among People Born Outside the U.S. by Gender – New York City, 2019-2024




Men consistently experienced the highest number of new HIV diagnoses, representing 72% of new diagnoses among people born outside the U.S. in 2024. This is slightly lower than the citywide proportion of diagnoses among men of 75% in 2024. The number of people newly diagnosed with HIV who were born outside the U.S. increased among men, women, and transgender women from 2019 to 2024. The number of new HIV diagnoses in all other gender groups remained low and relatively stable.

Number of New HIV Diagnoses and Percent Change Among People Born Outside the U.S. by Gender – New York City in 2019, 2023, and 2024




From 2019 to 2024, the number of people newly diagnosed with HIV who were born outside the U.S. increased among men (13%), women (52%), and transgender women (92%). More modest increases or decreases occurred for these groups from 2023 to 2024. All other gender groups experienced either no new diagnoses or relatively small numbers during these years; percent change calculations for these groups should be interpreted with caution.

Number of New HIV Diagnoses Among People Born Outside the U.S. by Race and Ethnicity – New York City, 2019-2024




Latino people consistently experienced the highest number of new HIV diagnoses, representing 55% of new diagnoses among people born outside the U.S. in 2024. This is higher than the citywide proportion of diagnoses among Latino people of 41% in 2024. From 2019 to 2024, the number of people newly diagnosed with HIV who were born outside the U.S. increased among Latino people and Black people. The number of new HIV diagnoses in all other race and ethnicity groups remained relatively stable.

Number of New HIV Diagnoses and Percent Change Among People Born Outside the U.S. by Race and Ethnicity – New York City in 2019, 2023, and 2024




From 2019 to 2024, the number of people newly diagnosed with HIV who were born outside the U.S. increased among Latino people (30%) and Black people (29%). Black people experienced a similar increase from 2023 to 2024, while Latino people experienced a decrease in the number of people newly diagnosed with HIV. All other groups experienced relatively low numbers of people newly diagnosed with HIV.

Number of New HIV Diagnoses Among People Born Outside the U.S. by Age Group – New York City, 2019-2024



People ages 20 to 39 years consistently experienced the highest number of new HIV diagnoses, representing a combined 65% of new diagnoses among people born outside the U.S. in 2024. This is similar to the citywide proportion of diagnoses among people ages 20 to 39 of 66% in 2024. From 2019 to 2024, the number of people newly diagnosed with HIV who were born outside the U.S. increased among people ages 20 to 29, 30 to 39, 40 to 49, and 60 and over. The number of new HIV diagnoses in all other age groups decreased or remained stable.

Number of New HIV Diagnoses and Percent Change Among People Born Outside the U.S. by Age Group – New York City in 2019, 2023, and 2024




From 2019 to 2024, the number of people newly diagnosed with HIV who were born outside the U.S. increased among people ages 20 to 29 (21%), 30 to 39 (33%), 40 to 49 (36%), and 60 and over (21%). From 2023 to 2024, more moderate increases or decreases were seen among all age groups except for people ages 60 and over, which increased by 53%.

Number of New HIV Diagnoses Among People Born Outside the U.S. by Race and Ethnicity and Age Group – New York City, 2024




Latino people born outside the U.S. ages 20 to 39 experienced the highest number of new HIV diagnoses, representing a combined 41% of new diagnoses among people born outside the U.S. in 2024.

Number of New HIV Diagnoses Among People Born Outside the U.S. by Borough of Residence – New York City, 2019-2024




People residing in Brooklyn experienced the highest number of new HIV diagnoses, representing 28% of new diagnoses among people born outside the U.S. in 2024. This is the same as the citywide proportion of diagnoses among people residing in Brooklyn. From 2019 to 2024, the number of people newly diagnosed with HIV who were born outside the U.S. increased among people residing in Brooklyn, Queens, the Bronx, and Staten Island. The number of new HIV diagnoses in all other borough of residence groups remained relatively stable.

Number of New HIV Diagnoses and Percent Change Among People Born Outside the U.S. by Borough of Residence – New York City in 2019, 2023, and 2024



From 2019 to 2024, the number of people newly diagnosed with HIV who were born outside the U.S. increased among people residing in Brooklyn (51%), Queens (17%), the Bronx (39%), and Staten Island (78%). From 2023 to 2024, more modest increases or decreases were seen among residents of all boroughs except for Brooklyn, which increased by 39%.

Number of New HIV Diagnoses Among People Born Outside the U.S. by Neighborhood Poverty Level¹ – New York City, 2019-2024



Medium-poverty neighborhoods consistently experienced the highest number of new HIV diagnoses, representing 47% of new diagnoses among people born outside the U.S. in 2024. This is higher than the citywide proportion of diagnoses among people residing in medium-poverty neighborhoods of 41% in 2024. From 2019 to 2024, the number of people newly diagnosed with HIV who were born outside the U.S. increased among people residing in medium-poverty neighborhoods, high-poverty neighborhoods, and very-high-poverty neighborhoods. The number of new HIV diagnoses in all other neighborhood poverty level groups remained relatively stable.




¹Neighborhood poverty level is determined by the proportion of residents living below the federal poverty level (FPL) in the NYC ZIP code of residence at diagnosis.

Low poverty=<10% below FPL; Medium poverty=10 to <20% below FPL; High poverty=20 to <30% below FPL; Very high poverty= \geq 30% below FPL.


As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Number of New HIV Diagnoses and Percent Change Among People Born Outside the U.S. by Neighborhood Poverty Level – New York City in 2019, 2023, and 2024




From 2019 to 2024, the number of people newly diagnosed with HIV who were born outside the U.S. increased among people residing in medium-poverty neighborhoods (42%), high-poverty neighborhoods (23%), and very-high-poverty neighborhoods (49%). All neighborhood poverty level groups experienced more modest increases or decreases from 2023 to 2024.

Number of New HIV Diagnoses Among People Born Outside the U.S. by Transmission Category – New York City, 2019-2024



Men who have sex with men consistently experienced the highest number of new HIV diagnoses, representing 61% of new diagnoses among people born outside the U.S. for whom data on transmission category were available in 2024. This is lower than the citywide proportion of diagnoses with transmission data among men who have sex with men of 65% in 2024. From 2019 to 2024, the number of people newly diagnosed with HIV who were born outside the U.S. increased among people with an unknown transmission category, people with heterosexual contact, and transgender people with sexual contact. The number of new HIV diagnoses in all other transmission category groups remained relatively stable or decreased.

Number of New HIV Diagnoses and Percent Change Among People Born Outside the U.S. by Transmission Category – New York City in 2019, 2023, and 2024




From 2019 to 2024, the number of people newly diagnosed with HIV who were born outside the U.S. increased among people with an unknown transmission category (83%), people with heterosexual contact (27%), and transgender people with sexual contact (100%). From 2023 to 2024, more modest increases or decreases were seen among all transmission category groups except for people with an unknown transmission category, which increased by 36%. Due to the relatively large number of people with an unknown transmission category, percent change calculations for all other groups should be interpreted with caution.

Care Outcomes Among People Newly Diagnosed With HIV Who Were Born Outside the United States


New York City

Initiation of Care¹ Within 30 Days of Diagnosis Among People Born Outside the U.S. – New York City, 2019–2024



From 2019 to 2024, initiation of care within 30 days of diagnosis among people born outside the U.S. increased four percentage points and was higher than New York City overall.

Initiation of Care¹ Within 30 Days of Diagnosis Among People Born Outside the U.S. by Demographic Group – New York City, 2024



*Data should be interpreted with caution because of small population size.

MSM-IDU=Men who have sex with men and inject drugs; TG-SC=Transgender people with sexual contact.


¹Initiation of care is defined as receiving a CD4, viral load, or genotype test after an HIV diagnosis. People diagnosed at death have been excluded.

²Neighborhood poverty level is determined by the proportion of residents living below the federal poverty level (FPL) in the NYC ZIP code of residence at diagnosis.

Low poverty=<10% below FPL; Medium poverty=10 to <20% below FPL; High poverty=20 to <30% below FPL; Very high poverty=>30% below FPL.


As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Initiation of Care¹ Within 30 Days of Diagnosis Among People Born Outside the U.S. by United Hospital Fund Neighborhood – New York City, 2024




The neighborhoods with the lowest proportions of initiation of care within 30 days of diagnosis among people born outside the U.S. were Kingsbridge – Riverdale (67%), Gramercy Park – Murray Hill (67%), and Bedford Stuyvesant – Crown Heights (72%).

Viral Suppression¹ Within Three Months of Diagnosis Among People Born Outside the U.S. – New York City, 2019-2024



From 2019 to 2024, viral suppression within three months of an HIV diagnosis among people born outside of the U.S. increased by seven percentage points and was predominately higher than New York City overall.

Viral Suppression¹ Within Three Months of Diagnosis Among People Born Outside the U.S. by Demographic Group – New York City, 2024



Differences in viral suppression within three months of an HIV diagnosis exist across demographic groups among people born outside the U.S.

*Data should be interpreted with caution because of small population size.

MSM-IDU=Men who have sex with men and inject drugs; TG-SC=Transgender people with sexual contact.


¹Viral suppression is defined as an HIV viral load in the calendar year <200 copies/mL within three months of diagnosis. People diagnosed at death have been excluded.

²Neighborhood poverty level is determined by the proportion of residents living below the federal poverty level (FPL) in the NYC ZIP code of residence at diagnosis.

Low poverty=<10% below FPL; Medium poverty=10 to <20% below FPL; High poverty=20 to <30% below FPL; Very high poverty=>30% below FPL.

As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Viral Suppression¹ Within Three Months of Diagnosis Among People Born Outside the U.S. by United Hospital Fund Neighborhood – New York City, 2024




The neighborhoods with the lowest proportions of people born outside the U.S. virally suppressed within three months of an HIV diagnosis were Pelham – Throgs Neck (47%), Southwest Queens (47%), Gramercy Park – Murray Hill (50%), and Ridgewood – Forest Hills (50%).

Care Outcomes Among People With HIV Who Were Born Outside the United States


New York City

Viral Suppression¹ Among People Born Outside the U.S. and Diagnosed With HIV² – New York City, 2019-2024



From 2019 to 2024, viral suppression among people born outside the U.S. with diagnosed HIV increased three percentage points and was higher than New York City overall.

Viral Suppression¹ Among People Born Outside the U.S. and Diagnosed With HIV² by Demographic Group – New York City, 2024



*Data should be interpreted with caution because of small population size.

MSM-IDU=Men who have sex with men and inject drugs; TG-SC=Transgender people with sexual contact.

¹Viral suppression is defined as the last HIV viral load in the calendar year <200 copies/mL. People diagnosed at death have been excluded.


²People diagnosed with HIV and viral suppression were calculated using the statistical weighting method. For more details and references, see Technical Notes.

³Neighborhood poverty level is determined by the proportion of residents living below the federal poverty level (FPL) in the NYC ZIP code of residence at diagnosis.

Low poverty=<10% below FPL; Medium poverty=10 to <20% below FPL; High poverty=20 to <30% below FPL; Very high poverty=>30% below FPL.

As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.


Viral Suppression¹ Among People Born Outside the U.S. and Diagnosed With HIV² by United Hospital Fund Neighborhood – New York City, 2024



The neighborhoods with the lowest proportions of virally suppressed people born outside of the U.S. and diagnosed with HIV were Bayside – Little Neck (75%) and Willowbrook (81%).

Proportion of People With HIV Born Outside the U.S. in Stages of the HIV Care Continuum^{1,2}

Overall and by Race and Ethnicity³ – New York City, 2024




In New York City, of approximately 23,800 people with HIV born outside the U.S., 84% had a suppressed viral load in 2024. This was higher than the citywide proportion of people with a suppressed viral load of 81%. There were differences in the HIV care continuum by race and ethnicity in 2024.

¹The HIV care continuum is a series of key stages for people with HIV. The denominator for each displayed proportion is the estimated number of people with HIV within a given group. For definitions of the stages of the continuum of care, see Technical Notes.

²Proportions in the care continuum may not align between stages due to the use of multiple data sources in calculations (e.g., proportion prescribed ART may be lower than the proportion virally suppressed)

³The estimated number of people with HIV by race and ethnicity may not sum to the overall value due to rounding and the use of specific estimated proportions of people with HIV who have been diagnosed within each race and ethnicity group.

Age-Adjusted¹ Death Rate per 1,000 People Born Outside the U.S. and Diagnosed With HIV – New York City, 2019–2024




The age-adjusted death rate declined by 13% since 2019 and 36% since its recent peak in 2020. The death rate among people born outside the U.S. was lower than the New York City overall rate.

Age-Adjusted¹ Death Rate per 1,000 People Born Outside the U.S. and Diagnosed With HIV by Demographic Group – New York City, 2024


6.8 deaths per 1,000 people with HIV born outside the U.S. overall




Race and ethnicity



6.8 deaths per 1,000 people with HIV born outside the U.S. overall



Neighborhood poverty²



Differences in the age-adjusted death rate exist across demographic groups among people born outside the U.S.

*Data should be interpreted with caution because of small population size.


¹Age-adjusted to the standard 2000 U.S. population. People newly diagnosed with HIV at death were excluded from the numerator.

²Neighborhood poverty level is determined by the proportion of residents living below the federal poverty level (FPL) in the NYC ZIP code of residence at diagnosis.

Low poverty=<10% below FPL; Medium poverty=10 to <20% below FPL; High poverty=20 to <30% below FPL; Very high poverty=>30% below FPL.


As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Age-Adjusted¹ Death Rate per 1,000 People Born Outside the U.S. and Diagnosed With HIV by United Hospital Fund Neighborhood – New York City, 2024



The neighborhoods with the highest age-adjusted death rates among people born outside the U.S. with HIV were Stapleton – St. George (36.8 per 1,000), Jamaica (28.7 per 1,000), and Flushing – Clearview (12.7 per 1,000).

Proportion of Deaths Among People Born Outside the U.S. With HIV by Cause of Death – New York City, 2023¹



In 2023, 79% of deaths among people with HIV were due to non-HIV-related causes. Among these, the top causes were cardiovascular disease (28%), non-HIV-related cancers (26%), and accidents (13%).

Appendix: How to Find Our Data

The New York City Department of Health and Mental Hygiene (NYC Health Department) issues the various publications related to our HIV surveillance data, including:



- **Annual HIV surveillance reports, surveillance slide sets, and statistics tables**, available at: <https://www.nyc.gov/site/doh/data/data-sets/hiv-aids-surveillance-and-epidemiology-reports.page>
- **HIV Care Status Reports**, available at: <https://www.nyc.gov/site/doh/health/health-topics/aids-hiv-care-status-reports-system.page>
- **HIV Care Continuum Dashboards**, available at: <https://www.nyc.gov/site/doh/health/health-topics/care-continuum-dashboard.page>

For HIV surveillance data requests, email HIVReport@health.nyc.gov. Please allow a minimum of two weeks for requests to be completed.

Appendix: Technical Notes

- **Inclusion criteria:** NYC HIV surveillance data include all people who are diagnosed with HIV by a provider located in NYC, regardless of their place of residence. NYC HIV surveillance investigates all people who were previously unknown to the NYC HIV Surveillance Registry. People who have an indication of previous HIV diagnosis, through health record review, interview, or federal duplication efforts (e.g., Routine Interstate Duplicate Review [RIDR]), are not included under people newly diagnosed with HIV in this report.
- **HIV Incidence:** HIV incidence is the number of people who acquired HIV in a population in a given period (such as a calendar year) as estimated based on a CD4 depletion model.¹ This differs from HIV diagnoses, which is the number of people who were newly diagnosed with HIV in a population in a given period (such as a calendar year), regardless of when they may actually have acquired HIV, which may have been many years prior to their diagnosis.
- **Gender Identity:** NYC HIV surveillance has routinely collected information gender identity since 2005 for newly reported cases. This report displays the following gender categories: men, women, transgender women, transgender men, and additional gender identities. In this report, people whose current gender identity differs from their sex assigned at birth are considered transgender people, and people who reported a nonbinary, genderqueer, gender nonconforming or any gender identity not previously listed are grouped under additional gender identities. Gender identities listed here are included without any intended hierarchy or prioritization – and are based on limited data reported to HIV surveillance. Classifying gender in surveillance requires accurate collection of sex assigned at birth and gender identity. Sex assigned at birth and gender information are collected from people's self-reports, their health care providers, or medical chart reviews. This information may or may not be complete or reflect self-identification. Reported numbers in this report among transgender people and people with additional gender identities are likely to be underestimates.
- **Race and Ethnicity:** NYC HIV surveillance collects data on race and ethnicity from multiple sources, including medical charts, provider reporting, vital statistics records, and patient interviews. Black, white, Asian or Pacific Islander, Native American, and multiracial race categories exclude Latino ethnicity. People with the ethnicity Latino are grouped in the Latino race and ethnicity category, regardless of their race classification. People not identified as Latino who identify with more than one race are classified under multiracial.
- **Area-Based Poverty:** Area-based poverty is based on NYC ZIP code of residence and is defined as the percentage of the population in a ZIP code with a household income that is below the federal poverty level. In this report, for HIV and AIDS diagnoses, ZIP code of residence at diagnosis; for people with HIV and deaths, ZIP code of residence on most recent record available. This measure is not available for people missing a ZIP code or living outside NYC. Income data used in this report are from the five-year American Community Survey (ACS) estimates centered on the year of the numerator data (for example, 2019 to 2023 ACS five-year estimate for 2021 data); if the preferred five-year file was not available, the most recent five-year ACS file was used. Cut points for area-based poverty categories in NYC were defined by a NYC Health Department work group.²



¹Source: NYC HIV Surveillance Registry; method: Song R, Hall HI, Green TA, Szwarcwald CL, Pantazis N. Using CD4 data to estimate HIV incidence, prevalence, and percent of undiagnosed infections in the United States. *J Acquir Immune Defic Syndr*. 2017;74(1):3-9. doi: 10.1097/QAI.0000000000001151 ²Toprani A, Hadler JL. Selecting and applying a standard area-based socioeconomic status measure for public health data: analysis for New York City. NYC Dept of Health and Mental Hygiene: Epi Research Report; May 2013.

Appendix: Technical Notes

- **Transmission Category:** NYC HIV surveillance collects data on behaviors possibly related to HIV transmission that occurred any time prior to diagnosis. Transmission categories include men who have sex with men, injection drug use, men who have sex with men and inject drugs, heterosexual contact, transgender people with sexual contact, perinatal transmission, and other. Men who have sex with men includes men with reported sexual contact with another man, and men with a history of a rectal sexually transmitted infection or proctitis and no other definitive transmission category. Injection drug use includes people with a history of taking nonprescribed drugs by injection, intravenously, intramuscularly or subcutaneously, excluding men reporting a history of sex with men. Men who have sex with men and inject drugs includes people meeting the definition of both the men who have sex with men and injection drug use categories as described above. Heterosexual contact includes people who had heterosexual sex with a person they know to have HIV, a person they know to have injected drugs, or a person they know to have received blood products. For women only, it also includes history of sex work, multiple sex partners, sexually transmitted infection, crack or cocaine use, sex with a bisexual man, probable heterosexual transmission as noted in a medical chart, or sex with a man and no injection drug use history. Transgender people with sexual contact includes people identified as transgender at any time who have reported sexual contact and no injection drug use history. Transgender people with injection drug use history are categorized under injection drug use history. Perinatal includes people who were exposed to HIV during gestation, birth or postpartum through breastfeeding to a parent with HIV. Other includes people who received treatment for hemophilia, people who received a transfusion or transplant, people with other health care-associated transmission, and children with non-perinatal transmission. Unknown includes people for whom data are not available to classify them in one of the transmission categories described above.
- **Death Data:** NYC HIV surveillance collects data on deaths among people with HIV occurring in NYC through matches with the NYC Vital Statistics registry, medical chart reviews, and provider reports, including on autopsies of people with HIV by the NYC Office of Chief Medical Examiner. Data on deaths occurring outside NYC are from matches with the U.S. Social Security Administration's Death Master File and CDC's National Death Index. At the time of publication of this report, death data for the reporting period are incomplete. They include preliminary NYC death data, National Death Index data, and partial Death Master File data.
- **Cause of Death:** In this report, cause of death is a person's underlying cause of death. For deaths occurring between 1984 and 1986, ICD-9 code 279.1 was used to denote AIDS-related deaths. For deaths occurring between 1987 and 1998, ICD-9 codes 042-044 were used to denote HIV- or AIDS-related deaths. For deaths occurring between 1999 and the most recent year, ICD-10 codes B20-B24 were used to denote HIV/AIDS-related deaths. For technical notes on cause of death by the NYC Health Department's Bureau of Vital Statistics, see nyc.gov/assets/doh/downloads/pdf/vs/2022sum.pdf. HIV infection and its management may contribute to causes of death classified as non-HIV-related, such as cardiovascular disease and certain cancers.^{1,2}

Appendix: Technical Notes on the HIV Care Continuum

NYC HIV Care Continuum: The care continuum is a common model used to quantify the progress of people with HIV through the stages of HIV care, with the ultimate goal being viral suppression. The stages of the care continuum are defined as follows:

- **People with HIV** is the estimated number of people diagnosed and undiagnosed with HIV. This estimate is calculated as the number of people diagnosed with HIV divided by the estimated proportion of people with HIV who are diagnosed, based on a CD4 depletion model.¹ All proportions in the Continuum use this number as the denominator.
- **HIV-Diagnosed** is calculated as the number of people with HIV retained in care plus the estimated number of people with HIV who were out of care, based on a statistical weighting method. This estimated number aims to account for migration out of NYC and therefore is different from the total number of people diagnosed and reported with HIV in NYC as seen in Figure 1 and Table 1 of the HIV surveillance annual report.²
- **Received care** is defined as people with HIV with one or more viral load or CD4 laboratory result reported in the calendar year to NYC HIV surveillance.³
- **Prescribed ART** is calculated as the number of people with HIV who received care multiplied by the estimated proportion of people with HIV prescribed ART in the previous 12 months, based on the proportion of NYC MMP participants whose medical record included documentation of ART prescription.⁴
- **Virally suppressed** is calculated as people with HIV in care with a most recent viral load measurement in the calendar year of less than 200 copies per milliliter, plus the estimated number of out-of-care people with HIV in the calendar year with a viral load of less than 200 copies per milliliter, based on a statistical weighting method.²

Appendix: Reporting HIV and AIDS Diagnoses for Health Care Providers

New York State (NYS) law requires health care providers to report HIV and AIDS diagnoses.

NYS Public Health Law¹ requires providers to report within seven days of diagnosis or receipt of laboratory results:

- New HIV diagnoses
- New AIDS diagnoses (if the patient has fewer than 200 CD4 cells per µL or an AIDS-related opportunistic infection)
- Previously diagnosed HIV or AIDS (if seeing the patient for the first time)

Providers must report within 24 hours of diagnosis:

- Acute HIV infections

Submit reports using the NYS Medical Provider HIV/AIDS and Partner/Contact Report Form (DOH-4189) by:

- Submitting the form electronically through the NYS Health Commerce System's Provider Portal at commerce.health.state.ny.us. For assistance with the portal, see the provider reporting guide at [Provider Reporting Guide](#) or call the NYS Department of Health at 518-474-4284.
- Obtaining paper forms from the NYC Health Department and arranging for the pickup of completed paper forms by calling 212-442-3388. You may also fax the completed form to the NYC Health Department at 347-396-8816. To protect patient confidentiality, completed forms must not be mailed to the NYC Health Department.

Providers should notify their patients newly diagnosed with HIV that they may be contacted by NYC Health Department's Assess.Connect.Engage. (ACE) Team who can assist them and their partners to:

- Assess health care and supportive service needs
- Connect patients who did not receive their HIV test results or missed their first medical appointment to HIV care
- Engage patient's partners in HIV testing, prevention, treatment, and supportive services, as needed

To contact the ACE Team, call 347-396-7601 Monday to Friday from 9 a.m. to 5 p.m. or email ACE@health.nyc.gov.



For more information and resources on reporting HIV and AIDS diagnoses, scan the QR code or visit: nyc.gov/health/hivprovider/reporting

Appendix: Acknowledgements

This report was prepared by the HIV Epidemiology Program in the NYC Health Department's Bureau of Hepatitis, HIV, and Sexually Transmitted Infections. We would like to acknowledge staff in the HIV Epidemiology Program's Surveillance Unit, ACE Team, Core HIV Surveillance Special Projects, and Data Support Unit, whose work is the foundation of this report.

The HIV Epidemiology Program's work depends on the participation of NYC providers, New Yorkers with HIV, community members and multiple other contributors. To them we are immensely indebted. Thank you.