

HIV Among Women – New York City, 2024

HIV Epidemiology Program
New York City Department of Health and Mental Hygiene
Published December 2025
nyc.gov/hivreports



Table of Contents

Description	Slide
HIV – New York City	
by gender	
number of women newly diagnosed	5
number of women newly diagnosed and percent change	6
Basic statistics of HIV among women – New York City	7
HIV among women – New York City	
number of women newly diagnosed	8
number of women newly diagnosed and percent change	9
rate of new diagnoses per 100,000 women	10
by race and ethnicity	
number of women newly diagnosed	11
number of women newly diagnosed and percent change	12
proportion of women newly diagnosed and all women in New York City	13
rate of new diagnoses per 100,000 women	14
by age group	
number of women newly diagnosed	15
number of women newly diagnosed and percent change	16
proportion of women newly diagnosed and all women in New York City	17
rate of new diagnoses per 100,000 women	18
by race and ethnicity and age group	
number of women newly diagnosed	19
by borough	
number of women newly diagnosed	20
number of women newly diagnosed and percent change	21
proportion of women newly diagnosed and all women in New York City	22
rate of new diagnoses per 100,000 women	23


Table of Contents

Description	Slide
HIV among women – New York City (continued)	
by neighborhood poverty level	
number of women newly diagnosed	24
number of women newly diagnosed and percent change	25
proportion of women newly diagnosed and all women in New York City	26
rate of new diagnoses per 100,000 women	27
by transmission category	
number of women newly diagnosed	28
number of women newly diagnosed and percent change	29
by place of birth	
number of women newly diagnosed	30
number of women newly diagnosed and percent change	31
Initiation of care within 30 days of diagnosis among women – New York City	33
by demographic group	34
by United Hospital Fund Neighborhood	35
Viral suppression within three months of diagnosis among women – New York City	36
by demographic group	37
by United Hospital Fund Neighborhood	38

Table of Contents


Description	Slide
Viral suppression among women diagnosed with HIV – New York City	40
by demographic group	41
by United Hospital Fund Neighborhood	42
HIV care continuum among women – New York City	
overall and by race and ethnicity	43
Age-adjusted death rate per 1,000 women with HIV – New York City	44
by demographic group	45
by United Hospital Fund Neighborhood	46
Deaths among women with HIV by cause – New York City	47
Appendices	
How to find our data	48
Technical notes	49
Technical notes on the HIV care continuum	51
Reporting HIV and AIDS diagnoses for health care providers	52
Acknowledgements	53

Number of People Newly Diagnosed With HIV by Gender – New York City, 2019-2024



Men consistently experienced the highest number of new HIV diagnoses, accounting for 75% of new diagnoses in 2024. The number of people newly diagnosed with HIV increased among women and among transgender women from 2019 to 2024. All other gender groups experienced decreases or relatively stable numbers of new HIV diagnoses. In 2024, women and transgender women represented a combined 25% of new diagnoses overall.

Number of People Newly Diagnosed With HIV and Percent Change by Gender – New York City in 2019, 2023, and 2024



The number of men newly diagnosed with HIV decreased 4% from 2019 to 2024 and was relatively stable from 2023 to 2024. From 2019 to 2024, the number of new HIV diagnoses increased among women (11%) and transgender women (30%); similar increases occurred among these groups from 2023 to 2024. All other gender groups experienced relatively small numbers of new diagnoses during these years; percent change calculations for these groups should be interpreted with caution.

Basic Statistics of HIV Among Women¹ – New York City, 2024

- **439 women newly diagnosed with HIV**
 - Including 75 women concurrently diagnosed with AIDS (17.1% of diagnoses)
- **331 women newly diagnosed with AIDS²**
- **There are an estimated 25,800 women with HIV³**
- **449 deaths among women with HIV**
 - 7.9 deaths per 1,000 women with HIV⁴

¹Women includes transgender women.


²Includes people concurrently diagnosed with HIV and AIDS.

³Approximate value calculated as the number of people with HIV divided by the estimated proportion of people with HIV who had been diagnosed, see Technical Notes for more details.

⁴Age-adjusted to the 2000 U.S. Standard Population. People newly diagnosed with HIV at death were excluded from the analysis. Death data for 2024 are incomplete.


As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Number of New HIV Diagnoses Among Women¹ – New York City, 2019-2024




**The number of women newly diagnosed with HIV increased from 2019 to 2024.
In 2024, women represented 25% of new diagnoses overall in New York City.**

Number of New HIV Diagnoses and Percent Change Among Women¹ – New York City in 2019, 2023, and 2024




The number of women newly diagnosed with HIV increased from 2019 to 2024 (14%) and from 2023 to 2024 (20%).

Rate of New HIV Diagnoses¹ per 100,000 Women² – New York City, 2019-2024




The rate of new HIV diagnoses among women increased by 26% from 2019 to 2024.

Number of New HIV Diagnoses Among Women¹ by Race and Ethnicity – New York City, 2019-2024




Black and Latina women consistently experienced the highest number of new HIV diagnoses, representing a combined 91% of new diagnoses among women in 2024. This is higher than the citywide proportion of diagnoses among Black and Latino people of 85% in 2024. From 2019 to 2024, the number of women newly diagnosed with HIV increased among Latina women. Black women experienced a steep decline from 2019 to 2020, and a subsequent increase from 2020 to 2024. The number of new HIV diagnoses in all other race and ethnicity groups decreased or remained relatively stable.

Number of New HIV Diagnoses and Percent Change Among Women¹ by Race and Ethnicity – New York City in 2019, 2023, and 2024




From 2019 to 2024, the number of women newly diagnosed with HIV increased among Latina women (46%). All other race and ethnicity groups experience relatively low numbers of people newly diagnosed with HIV. From 2023 to 2024, increases occurred among Latina women (15%) and Black women (24%).

Proportion of Women¹ Newly Diagnosed With HIV and All Women^{2,3} by Race and Ethnicity – New York City, 2024




The proportions of new HIV diagnoses among Black and Latina women were higher than those groups' proportions in the New York City population.

Rate of New HIV Diagnoses^{1,2} per 100,000 Women³ by Race and Ethnicity – New York City, 2019-2024




Black women consistently experienced the highest rate of new HIV diagnoses. From 2019 to 2024, the rate of new HIV diagnoses per 100,000 women increased among Latina women (73%) and among Black women (7%) from 2019 to 2024. The rate of new HIV diagnoses among Native American women fluctuated; the count remained low, and the rate should be interpreted with caution. The rate of new HIV diagnoses in all other race and ethnicity groups decreased or remained relatively stable.

Number of New HIV Diagnoses Among Women¹ by Age Group – New York City, 2019-2024




Women ages 20 to 39 years consistently experienced the highest number of new HIV diagnoses, representing a combined 55% of new diagnoses among women in 2024. This is lower than the citywide proportion of diagnoses among people ages 20 to 39 of 66% in 2024. From 2019 to 2024, the number of women newly diagnosed with HIV increased among women ages 20 to 29, 30 to 39, 40 to 49, and 60 and over. The number of women newly diagnosed with HIV in all other age groups decreased or remained relatively stable.

Number of New HIV Diagnoses and Percent Change Among Women¹ by Age Group – New York City in 2019, 2023, and 2024




From 2019 to 2024, the number of women newly diagnosed with HIV increased among women ages 20 to 29 (24%), 30 to 39 (18%), 40 to 49 (20%), and 60 and over (48%). From 2023 to 2024, all age groups except girls ages 0 to 12 experienced an increase, with the largest percent increase among women ages 60 and over (58%).

Proportion of Women¹ Newly Diagnosed With HIV and All Women^{2,3} by Age Group – New York City, 2024



The proportions of new HIV diagnoses among women ages 20 to 39 were higher than those groups' proportions in the New York City population.

Rate of New HIV Diagnoses^{1,2} per 100,000 Women³ by Age Group – New York City, 2019-2024




Women ages 20 to 39 consistently experienced the highest rates of new HIV diagnoses. From 2019 to 2024, the rate of new HIV diagnoses per 100,000 women increased among women ages 20 to 29 (48%), 30 to 39 (31%), and 40 to 49 (32%). The rate of new HIV diagnoses in all other age groups decreased or remained relatively stable.

Number of New HIV Diagnoses Among Women¹ by Race and Ethnicity and Age Group – New York City, 2024




Black and Latina women ages 20 to 39 experienced the highest number of new HIV diagnoses, representing a combined 51% of new diagnoses among women in 2024.

Number of New HIV Diagnoses Among Women¹ by Borough of Residence – New York City, 2019-2024




Women residing in Brooklyn and the Bronx consistently experienced the highest number of new HIV diagnoses, representing a combined 55% of new diagnoses in 2024. This is higher than the citywide proportion of diagnoses among people residing in Brooklyn of 50% in 2024. From 2019 to 2024, the number of women newly diagnosed with HIV increased among women residing in Brooklyn, Queens, and Manhattan. The number of women newly diagnosed with HIV in all other borough of residence groups decreased or remained relatively stable.

Number of New HIV Diagnoses and Percent Change Among Women¹ by Borough of Residence – New York City in 2019, 2023, and 2024




From 2019 to 2024, the number of women newly diagnosed with HIV increased among women residing in Brooklyn (36%), Queens (19%), and Manhattan (48%). From 2023 to 2024, people residing in these same boroughs experienced increases. The number of women newly diagnosed with HIV decreased or remained relatively stable in all other borough of residence groups.

Proportion of Women¹ Newly Diagnosed With HIV and All Women^{2,3} by Borough of Residence³ – New York City, 2024



The proportions of new HIV diagnoses among women residing in the Bronx and Brooklyn were higher than those groups' proportions in the New York City population.

Rate of New HIV Diagnoses^{1,2} per 100,000 Women³ by Borough of Residence – New York City, 2019-2024



Women in the Bronx consistently experienced the highest rate of new HIV diagnoses. From 2019 to 2024, the rate of new HIV diagnoses per 100,000 women increased among women residing in Brooklyn (46%), Manhattan (60%), and Queens (27%). The rate of new HIV diagnoses in all other borough of residence groups either decreased or remained relatively stable.




¹Rates are a measure that account for population size, allowing for a clearer comparison in new HIV diagnoses. Rates were calculated using Health Department population estimates, modified from U.S. Census Bureau intercensal population estimates.

²Excludes people newly diagnosed with HIV in New York City who were residing outside of New York City at the time of diagnosis.


³Women includes transgender women.

Number of New HIV Diagnoses Among Women¹ by Neighborhood Poverty Level² – New York City, 2019-2024



Women residing in medium-poverty neighborhoods consistently experienced the highest number of new HIV diagnoses, representing 38% of new diagnoses among women in 2024. This is lower than the citywide proportion of diagnoses among people residing in medium-poverty neighborhoods of 41% in 2024. The number of women newly diagnosed with HIV increased among women residing in medium-poverty neighborhoods from 2019 to 2024. The number of new HIV diagnoses in all other neighborhood poverty level groups either decreased or remained relatively stable.

Number of New HIV Diagnoses and Percent Change Among Women¹ by Neighborhood Poverty Level² – New York City in 2019, 2023, and 2024



From 2019 to 2024, the number of women newly diagnosed with HIV increased among women residing in medium-poverty neighborhoods (39%). From 2023 to 2024, increases occurred among women residing in low-poverty neighborhoods (59%), medium-poverty neighborhoods (15%), high-poverty neighborhoods (25%), and very-high-poverty neighborhoods (24%).


¹Women includes transgender women.

²Neighborhood poverty level is determined by the proportion of residents living below the federal poverty level (FPL) in the NYC ZIP code of residence at diagnosis.
Low poverty=<10% below FPL; Medium poverty=10 to <20% below FPL; High poverty=20 to <30% below FPL; Very high poverty=>30% below FPL.

³One-year increases or decreases are not indicative of a trend and should be interpreted with caution.

As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Proportion of Women¹ Newly Diagnosed With HIV and All Women^{2,3} by Neighborhood Poverty Level^{4,5} – New York City, 2024



The proportions of new HIV among women residing in high- or very-high-poverty neighborhoods were higher than those groups' proportions in the New York City population.

¹Women includes transgender women.

²NYC population calculated using Health Department population estimates, modified from U.S. Census Bureau intercensal population estimates.

³Excludes people newly diagnosed with HIV in New York City who were residing outside of New York City at the time of diagnosis.


⁴Neighborhood poverty level is determined by the proportion of residents living below the federal poverty level (FPL) in the NYC ZIP code of residence at diagnosis.

Low poverty=<10% below FPL; Medium poverty=10 to <20% below FPL; High poverty=20 to <30% below FPL; Very high poverty=≥30% below FPL.

⁵Proportions exclude people living in neighborhoods with an unknown poverty level

As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Rate of New HIV Diagnoses^{1,2} per 100,000 Women³ by Neighborhood Poverty Level⁴ – New York City, 2019-2024



Women residing in very-high-poverty neighborhoods consistently experienced the highest rate of new HIV diagnoses. From 2019 to 2024, the rate of new HIV diagnoses per 100,000 women increased among women residing in very-high-poverty neighborhoods (22%), high-poverty neighborhoods (20%), and medium-poverty neighborhoods (50%).

¹Rates are a measure that account for population size, allowing for a clearer comparison in new HIV diagnoses. Rates were calculated using Health Department population estimates, modified from U.S. Census Bureau intercensal population estimates.

²Excludes people newly diagnosed with HIV in New York City who were residing outside of New York City at the time of diagnosis.


³Women includes transgender women.

⁴Neighborhood poverty level is determined by the proportion of residents living below the federal poverty level (FPL) in the NYC ZIP code of residence at diagnosis.

Low poverty=<10% below FPL; Medium poverty=10 to <20% below FPL; High poverty=20 to <30% below FPL; Very high poverty=>30% below FPL.


As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Number of New HIV Diagnoses Among Women¹ by Transmission Category – New York City, 2019-2024



Women with heterosexual contact consistently experienced the highest number of new HIV diagnoses, representing 77% of new diagnoses among women for whom data on transmission category were available in 2024. From 2019 to 2024, the number of women newly diagnosed with HIV increased among women with an unknown transmission category and among transgender women with sexual contact. The number of new HIV diagnoses in all other transmission category groups either decreased or remained relatively stable.


Number of New HIV Diagnoses and Percent Change Among Women¹ by Transmission Category – New York City in 2019, 2023, and 2024



From 2019 to 2024, the number of women newly diagnosed with HIV increased among women with an unknown transmission category (140%) and among transgender women with sexual contact (25%).


From 2023 to 2024, these groups also experienced increases. Due to the relatively large number of women with an unknown transmission category, percent change calculations for all other groups should be interpreted with caution.

Number of New HIV Diagnoses Among Women¹ by Place of Birth – New York City, 2019-2024



Women born outside of the United States experienced the highest number of new HIV diagnoses in 2024, representing 40% of new diagnoses among women. This is higher than the citywide proportion of diagnoses among people born outside the United States of 35% in 2024. From 2019 to 2024, the number of women newly diagnosed with HIV increased among women born outside of the United States and among women with an unknown place of birth. The number new HIV diagnoses in all other place of birth groups either decreased or remained relatively stable.

Number of New HIV Diagnoses and Percent Change Among Women¹ by Place of Birth – New York City in 2019, 2023, and 2024




From 2019 to 2024, the number of women newly diagnosed with HIV increased among women born outside of the United States (56%) and among women with an unknown place of birth (133%). From 2023 to 2024, these groups also experienced increases. Due to the relatively large number of women with an unknown place of birth, percent change calculations for all other groups should be interpreted with caution.

Care Outcomes Among Women Newly Diagnosed With HIV


New York City

Initiation of Care¹ Within 30 Days of Diagnosis Among Women² – New York City, 2019–2024



From 2019 to 2024, initiation of care within 30 days of diagnosis among women newly diagnosed with HIV increased by six percentage points and was approximately equal to New York City overall.

Initiation of Care¹ Within 30 Days of Diagnosis Among Women² by Demographic Group – New York City, 2024



Differences in initiation of care within 30 days of diagnosis exist across demographic groups among women.

*Data should be interpreted with caution because of small population size.

TG-SC=Transgender people with sexual contact.

¹Initiation of care is defined as receiving a CD4, viral load, or genotype test after an HIV diagnosis. People diagnosed at death have been excluded.


²Women includes transgender women.

³Neighborhood poverty level is determined by the proportion of residents living below the federal poverty level (FPL) in the NYC ZIP code of residence at diagnosis.

Low poverty=<10% below FPL; Medium poverty=10 to <20% below FPL; High poverty=20 to <30% below FPL; Very high poverty=>30% below FPL.


As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Initiation of Care¹ Within 30 Days of Diagnosis Among Women² by United Hospital Fund Neighborhood – New York City, 2024




The neighborhoods with the lowest proportions of initiation of care within 30 days of diagnosis among women were Coney Island – Sheepshead Bay (50%), Pelham – Throgs Neck (57%), Southeast Queens (67%), and Northeast Bronx (67%).

Viral Suppression¹ Within Three Months of Diagnosis Among Women² – New York City, 2019–2024



From 2019 to 2024, viral suppression within three months of an HIV diagnosis among women increased by seven percentage points and was approximately equal to New York City overall.

Viral Suppression¹ Within Three Months of Diagnosis Among Women² by Demographic Group – New York City, 2024



Differences exist in viral suppression within three months of an HIV diagnosis across demographic groups among women.

*Data should be interpreted with caution because of small population size.

TG-SC=Transgender people with sexual contact.

¹Viral suppression is defined as an HIV viral load in the calendar year <200 copies/mL within three months of diagnosis. People diagnosed at death have been excluded.


²Women includes transgender women.

³Neighborhood poverty level is determined by the proportion of residents living below the federal poverty level (FPL) in the NYC ZIP code of residence at diagnosis.

Low poverty=<10% below FPL; Medium poverty=10 to <20% below FPL; High poverty=20 to <30% below FPL; Very high poverty=>30% below FPL.

As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Viral Suppression¹ Within Three Months of Diagnosis Among Women² by United Hospital Fund Neighborhood – New York City, 2024




The neighborhoods with the lowest proportions of women virally suppressed within three months of an HIV diagnosis were Ridgewood – Forest Hills (33%), Stapleton – St. George (43%), and Pelham – Throgs Neck (43%).

Care Outcomes Among Women With HIV


New York City

Viral Suppression¹ Among Women² Diagnosed With HIV³ – New York City, 2019-2024



From 2019 to 2024, viral suppression among women diagnosed with HIV increased by three percentage points and was approximately equal to New York City overall.

Viral Suppression¹ Among Women² Diagnosed With HIV³ by Demographic Group – New York City, 2024



Differences in viral suppression exist across demographic group among women.

*Data should be interpreted with caution because of small population size.

TG-SC=Transgender people with sexual contact.

¹Viral suppression is defined as the last HIV viral load in the calendar year <200 copies/mL. People diagnosed at death have been excluded.

²Women includes transgender women.

³People diagnosed with HIV and viral suppression were calculated using the statistical weighting method. For more details and references, see Technical Notes.

⁴Neighborhood poverty level is determined by the proportion of residents living below the federal poverty level (FPL) in the NYC ZIP code of residence at diagnosis.


Low poverty=<10% below FPL; Medium poverty=10 to <20% below FPL; High poverty=20 to <30% below FPL; Very high poverty=>30% below FPL.

As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Viral Suppression¹ Among Women² Diagnosed With HIV³ by United Hospital Fund Neighborhood – New York City, 2024

Viral suppression among
people diagnosed with HIV
by UHF neighborhood


- 78% to 83%
- more than 83% to 86%
- more than 86% to 88%
- more than 88% to 95%
- Non-residential areas



The neighborhoods with the lowest proportions of virally suppressed women diagnosed with HIV were Central Harlem – Morningside Heights (78%), Downtown – Heights – Park Slope (79%), and Gramercy Park – Murray Hill (80%).

Proportion of Women¹ With HIV in Stages of the HIV Care Continuum^{2,3}

Overall and by Race and Ethnicity⁴ – New York City, 2024



Of approximately 25,800 women with HIV in 2024, 81% had a suppressed viral load. This was the same as the citywide proportion of people with a suppressed viral load. There were differences in the HIV care continuum among women by race and ethnicity in 2024.

¹Women includes transgender women.

²The HIV care continuum is a series of key stages for people with HIV. The denominator for each displayed proportion is the estimated number of people with HIV within a given group.


³Proportions in the care continuum may not align between stages due to the use of multiple data sources in calculations (e.g., proportion prescribed ART may be lower than the proportion virally suppressed)

⁴The estimated number of people with HIV by race and ethnicity may not sum to the overall value due to rounding and the use of specific estimated proportions of people with HIV who have been diagnosed within each race and ethnicity group.

For definitions of the stages of the continuum of care, see Technical Notes.

As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Age-Adjusted¹ Death Rate per 1,000 Women² With HIV – New York City, 2019-2024



The age-adjusted death rate among women with HIV declined by 14% since 2019 and 40% since its recent peak in 2020.
The death rate among women was approximately equal to New York City overall rate from 2019 to 2024

Age-Adjusted¹ Death Rate per 1,000 Women² With HIV by Demographic Group – New York City, 2024



Differences in the age-adjusted death rate exist across demographic groups among women.

*Data should be interpreted with caution because of small population size.

¹Age-adjusted to the standard 2000 U.S. population. People newly diagnosed with HIV at death were excluded from the numerator.


²Women includes transgender women.

³Neighborhood poverty level is determined by the proportion of residents living below the federal poverty level (FPL) in the NYC ZIP code of residence at diagnosis.

Low poverty=<10% below FPL; Medium poverty=10 to <20% below FPL; High poverty=20 to <30% below FPL; Very high poverty=>30% below FPL.


As reported to the New York City Department of Health and Mental Hygiene by March 31, 2025.

Age-Adjusted¹ Death Rate per 1,000 Women² With HIV by United Hospital Fund Neighborhood – New York City, 2024



The neighborhoods with the highest age-adjusted death rates among women with HIV were Jamaica (31.0 per 1,000), Greenpoint (14.6 per 1,000), and Stapleton – St. George (13.6 per 1,000).

Proportion of Deaths Among Women¹ With HIV by Cause of Death – New York City, 2023²



In 2023, 82% of deaths among women with HIV were due to non-HIV-related causes. Among these, the top causes were cardiovascular disease (33%), accidents (19%), and non-HIV-related cancers (19%).

¹Women includes transgender women.

²Cause of death data are not yet available for 2024.

³ICD10 codes B20-B24 were used to denote HIV-related deaths. For technical notes on cause of death by the NYC Health Department's Office of Vital Statistics, see nyc.gov/assets/doh/downloads/pdf/vs/2022sum.pdf.

Appendix: How to Find Our Data

The New York City Department of Health and Mental Hygiene (NYC Health Department) issues the various publications related to our HIV surveillance data, including:



- **Annual HIV surveillance reports, surveillance slide sets, and statistics tables**, available at: <https://www.nyc.gov/site/doh/data/data-sets/hiv-aids-surveillance-and-epidemiology-reports.page>
- **HIV Care Status Reports**, available at: <https://www.nyc.gov/site/doh/health/health-topics/aids-hiv-care-status-reports-system.page>
- **HIV Care Continuum Dashboards**, available at: <https://www.nyc.gov/site/doh/health/health-topics/care-continuum-dashboard.page>

For HIV surveillance data requests, email HIVReport@health.nyc.gov. Please allow a minimum of two weeks for requests to be completed.

Appendix: Technical Notes

- **Inclusion criteria:** NYC HIV surveillance data include all people who are diagnosed with HIV by a provider located in NYC, regardless of their place of residence. NYC HIV surveillance investigates all people who were previously unknown to the NYC HIV Surveillance Registry. People who have an indication of previous HIV diagnosis, through health record review, interview, or federal duplication efforts (e.g., Routine Interstate Duplicate Review [RIDR]), are not included under people newly diagnosed with HIV in this report.
- **HIV Incidence:** HIV incidence is the number of people who acquired HIV in a population in a given period (such as a calendar year) as estimated based on a CD4 depletion model.¹ This differs from HIV diagnoses, which is the number of people who were newly diagnosed with HIV in a population in a given period (such as a calendar year), regardless of when they may actually have acquired HIV, which may have been many years prior to their diagnosis.
- **Gender Identity:** NYC HIV surveillance has routinely collected information on gender identity since 2005 for newly reported cases. This report displays the following gender categories: men, women, transgender women, transgender men, and additional gender identities. In this report, people whose current gender identity differs from their sex assigned at birth are considered transgender people, and people who reported a nonbinary, genderqueer, gender nonconforming or any gender identity not previously listed are grouped under additional gender identities. Gender identities listed here are included without any intended hierarchy or prioritization – and are based on limited data reported to HIV surveillance. Classifying gender in surveillance requires accurate collection of sex assigned at birth and gender identity. Sex assigned at birth and gender information are collected from people's self-reports, their health care providers, or medical chart reviews. This information may or may not be complete or reflect self-identification. Reported numbers in this report among transgender people and people with additional gender identities are likely to be underestimates.
- **Race and Ethnicity:** NYC HIV surveillance collects data on race and ethnicity from multiple sources, including medical charts, provider reporting, vital statistics records, and patient interviews. Black, white, Asian or Pacific Islander, Native American, and multiracial race categories exclude Latino ethnicity. People with the ethnicity Latino are grouped in the Latino race and ethnicity category, regardless of their race classification. People not identified as Latino who identify with more than one race are classified under multiracial.
- **Area-Based Poverty:** Area-based poverty is based on NYC ZIP code of residence and is defined as the percentage of the population in a ZIP code with a household income that is below the federal poverty level. In this report, for HIV and AIDS diagnoses, ZIP code of residence at diagnosis; for people with HIV and deaths, ZIP code of residence on most recent record available. This measure is not available for people missing a ZIP code or living outside NYC. Income data used in this report are from the five-year American Community Survey (ACS) estimates centered on the year of the numerator data (for example, 2019 to 2023 ACS five-year estimate for 2021 data); if the preferred five-year file was not available, the most recent five-year ACS file was used. Cut points for area-based poverty categories in NYC were defined by a NYC Health Department work group.²



¹Source: NYC HIV Surveillance Registry; method: Song R, Hall HI, Green TA, Szwarcwald CL, Pantazis N. Using CD4 data to estimate HIV incidence, prevalence, and percent of undiagnosed infections in the United States. *J Acquir Immune Defic Syndr*. 2017;74(1):3-9. doi: 10.1097/QAI.0000000000001151 ²Toprani A, Hadler JL. Selecting and applying a standard area-based socioeconomic status measure for public health data: analysis for New York City. NYC Dept of Health and Mental Hygiene: Epi Research Report; May 2013.

Appendix: Technical Notes

- **Transmission Category:** NYC HIV surveillance collects data on behaviors possibly related to HIV transmission that occurred any time prior to diagnosis. Transmission categories include men who have sex with men, injection drug use, men who have sex with men and inject drugs, heterosexual contact, transgender people with sexual contact, perinatal transmission, and other. Men who have sex with men includes men with reported sexual contact with another man, and men with a history of a rectal sexually transmitted infection or proctitis and no other definitive transmission category. Injection drug use includes people with a history of taking nonprescribed drugs by injection, intravenously, intramuscularly or subcutaneously, excluding men reporting a history of sex with men. Men who have sex with men and inject drugs includes people meeting the definition of both the men who have sex with men and injection drug use categories as described above. Heterosexual contact includes people who had heterosexual sex with a person they know to have HIV, a person they know to have injected drugs, or a person they know to have received blood products. For women only, it also includes history of sex work, multiple sex partners, sexually transmitted infection, crack or cocaine use, sex with a bisexual man, probable heterosexual transmission as noted in a medical chart, or sex with a man and no injection drug use history. Transgender people with sexual contact includes people identified as transgender at any time who have reported sexual contact and no injection drug use history. Transgender people with injection drug use history are categorized under injection drug use history. Perinatal includes people who were exposed to HIV during gestation, birth or postpartum through breastfeeding to a parent with HIV. Other includes people who received treatment for hemophilia, people who received a transfusion or transplant, people with other health care-associated transmission, and children with non-perinatal transmission. Unknown includes people for whom data are not available to classify them in one of the transmission categories described above.
- **Death Data:** NYC HIV surveillance collects data on deaths among people with HIV occurring in NYC through matches with the NYC Vital Statistics registry, medical chart reviews, and provider reports, including on autopsies of people with HIV by the NYC Office of Chief Medical Examiner. Data on deaths occurring outside NYC are from matches with the U.S. Social Security Administration's Death Master File and CDC's National Death Index. At the time of publication of this report, death data for the reporting period are incomplete. They include preliminary NYC death data, National Death Index data, and partial Death Master File data.
- **Cause of Death:** In this report, cause of death is a person's underlying cause of death. For deaths occurring between 1984 and 1986, ICD-9 code 279.1 was used to denote AIDS-related deaths. For deaths occurring between 1987 and 1998, ICD-9 codes 042-044 were used to denote HIV- or AIDS-related deaths. For deaths occurring between 1999 and the most recent year, ICD-10 codes B20-B24 were used to denote HIV/AIDS-related deaths. For technical notes on cause of death by the NYC Health Department's Bureau of Vital Statistics, see nyc.gov/assets/doh/downloads/pdf/vs/2022sum.pdf. HIV infection and its management may contribute to causes of death classified as non-HIV-related, such as cardiovascular disease and certain cancers.^{1,2}

Appendix: Technical Notes on the HIV Care Continuum

NYC HIV Care Continuum: The care continuum is a common model used to quantify the progress of people with HIV through the stages of HIV care, with the ultimate goal being viral suppression. The stages of the care continuum are defined as follows:

- **People with HIV** is the estimated number of people diagnosed and undiagnosed with HIV. This estimate is calculated as the number of people diagnosed with HIV divided by the estimated proportion of people with HIV who are diagnosed, based on a CD4 depletion model.¹ All proportions in the Continuum use this number as the denominator.
- **HIV-Diagnosed** is calculated as the number of people with HIV retained in care plus the estimated number of people with HIV who were out of care, based on a statistical weighting method. This estimated number aims to account for migration out of NYC and therefore is different from the total number of people diagnosed and reported with HIV in NYC as seen in Figure 1 and Table 1 of the HIV surveillance annual report.²
- **Received care** is defined as people with HIV with one or more viral load or CD4 laboratory result reported in the calendar year to NYC HIV surveillance.³
- **Prescribed ART** is calculated as the number of people with HIV who received care multiplied by the estimated proportion of people with HIV prescribed ART in the previous 12 months, based on the proportion of NYC MMP participants whose medical record included documentation of ART prescription.⁴
- **Virally suppressed** is calculated as people with HIV in care with a most recent viral load measurement in the calendar year of less than 200 copies per milliliter, plus the estimated number of out-of-care people with HIV in the calendar year with a viral load of less than 200 copies per milliliter, based on a statistical weighting method.²

Appendix: Reporting HIV and AIDS Diagnoses for Health Care Providers

New York State (NYS) law requires health care providers to report HIV and AIDS diagnoses.

NYS Public Health Law¹ requires providers to report within seven days of diagnosis or receipt of laboratory results:

- New HIV diagnoses
- New AIDS diagnoses (if the patient has fewer than 200 CD4 cells per µL or an AIDS-related opportunistic infection)
- Previously diagnosed HIV or AIDS (if seeing the patient for the first time)

Providers must report within 24 hours of diagnosis:

- Acute HIV infections

Submit reports using the NYS Medical Provider HIV/AIDS and Partner/Contact Report Form (DOH-4189) by:

- Submitting the form electronically through the NYS Health Commerce System's Provider Portal at commerce.health.state.ny.us. For assistance with the portal, see the provider reporting guide at [Provider Reporting Guide](#) or call the NYS Department of Health at 518-474-4284.
- Obtaining paper forms from the NYC Health Department and arranging for the pickup of completed paper forms by calling 212-442-3388. You may also fax the completed form to the NYC Health Department at 347-396-8816. To protect patient confidentiality, completed forms must not be mailed to the NYC Health Department.

Providers should notify their patients newly diagnosed with HIV that they may be contacted by NYC Health Department's Assess.Connect.Engage. (ACE) Team who can assist them and their partners to:

- Assess health care and supportive service needs
- Connect patients who did not receive their HIV test results or missed their first medical appointment to HIV care
- Engage patient's partners in HIV testing, prevention, treatment, and supportive services, as needed

To contact the ACE Team, call 347-396-7601 Monday to Friday from 9 a.m. to 5 p.m. or email ACE@health.nyc.gov.



For more information and resources on reporting HIV and AIDS diagnoses, scan the QR code or visit: nyc.gov/health/hivprovider/reporting

Appendix: Acknowledgements

This report was prepared by the HIV Epidemiology Program in the NYC Health Department's Bureau of Hepatitis, HIV, and Sexually Transmitted Infections. We would like to acknowledge staff in the HIV Epidemiology Program's Surveillance Unit, ACE Team, Core HIV Surveillance Special Projects, and Data Support Unit, whose work is the foundation of this report.

The HIV Epidemiology Program's work depends on the participation of NYC providers, New Yorkers with HIV, community members and multiple other contributors. To them we are immensely indebted. Thank you.