



NEW YORK CITY DEPARTMENT OF
HEALTH AND MENTAL HYGIENE
Ashwin Vasani, MD, PhD
Commissioner

2022 Health Advisory #31: CDC Health Advisory: Interim Guidance for Clinicians to Prioritize Antiviral Treatment of Influenza in the Setting of Reduced Availability of Oseltamivir

December 16, 2022

Dear Colleagues,

Please see the **Center's for Disease Control and Prevention (CDC) Health Advisory: Interim Guidance for Clinicians to Prioritize Antiviral Treatment of Influenza in the Setting of Reduced Availability of Oseltamivir**, below.

Following are some key updates for New York City (NYC):

- Influenza is circulating at very high levels in NYC and influenza hospitalizations have increased significantly. For the week ending December 10, 2022, influenza-like illness made up 13% of all weekly emergency department, urgent care, and primary care clinic visit in NYC, and there were three reported influenza-associated pediatric deaths.
 - NYC weekly influenza reports are available [here](#). NYC influenza hospitalization data are included in the New York State weekly influenza activity report found [here](#).
- The NYC Commissioner of Health has issued an [Advisory](#), recommending mask use and other precautions, given the already high levels of COVID-19, influenza, and respiratory syncytial virus (RSV). Advise patients to take precautions to reduce their risk of illness, especially patients ages 65 and older and others at increased risk of complications.
- NYC pharmacies and hospitals are reporting spot shortages of generic oseltamivir.
- When there is limited availability of oseltamivir use clinical judgement to guide decision making and:
 - Prioritize oseltamivir treatment for patients with suspected or lab-diagnosed influenza who are hospitalized.
 - Prioritize oseltamivir for residents of congregate settings, such as long-term care facilities (LTCFs), who test positive for influenza, and for empiric treatment or chemoprophylaxis of residents during an influenza outbreak in the facility.
 - Among outpatients, prioritize antiviral treatment for patients who test positive for influenza who are [at increased risk of complications](#) such as older adults, very young children (age < 2 years), pregnant people, and people of all ages with certain chronic medical conditions. If oseltamivir is unavailable, oral baloxavir or inhaled zanamivir can be used for early treatment of outpatients at increased risk for complications who present with uncomplicated influenza, depending upon age and contraindications (see table).
- For patients with respiratory illness at high risk for complications, testing for both COVID-19 and influenza is recommended to guide antiviral treatment, preferably using a rapid molecular assay for flu when available.



- Because of the risk of co-infection, a positive test for SARS-CoV-2 does **not** exclude influenza infection, and vice versa.
- Vaccines remain the best way to protect against [influenza](#) and [COVID-19](#).
- Resources:
 - NYC Department of Health and Mental Hygiene: [City Health Information: Influenza Prevention and Control, 2022-2023](#)
 - CDC: [2022-2023 Seasonal Influenza Testing and Treatment During the COVID-19 Pandemic](#)
 - NYC Health Advisory #28: [Prescribe COVID-19 Therapeutics to Prevent Severe Disease, Hospitalization and Death This Winter](#)

Antiviral drug	Route of Administration	Recommended Ages for Treatment
Oseltamivir	Oral (twice daily x 5d)	All ages
Zanamivir	Inhaled (twice daily x 5d)	≥7 years
Peramivir	Intravenous (single infusion)	≥6 months
Baloxavir	Oral (single dose)	≥5 years (otherwise healthy) ≥12 years (high-risk)

Source: <https://www.cdc.gov/flu/professionals/antivirals/summary-clinicians.htm>

Distributed via the CDC Health Alert Network
December 14, 2022, 4:00 PM ET
CDCHAN-00482

Summary

Seasonal influenza activity is high across the United States. The Centers for Disease Control and Prevention (CDC) estimates that in the 2022-2023 season to date, there have been at least 13 million illnesses, 120,000 hospitalizations, and 7,300 deaths from influenza ([Weekly U.S. Influenza Surveillance Report | CDC](#)). While the [Food and Drug Administration](#) (FDA) has not indicated shortages of oseltamivir (generic or Tamiflu) in any of its forms (capsules, oral suspension), CDC has received numerous anecdotal reports of availability issues for generic oseltamivir in some locations [1]. This may continue to occur in some communities as influenza activity continues.

This Health Alert Network (HAN) Health Advisory provides clinicians and public health officials with guidance for prioritizing oseltamivir for treatment and information on other influenza antivirals that are recommended for treating influenza in areas where oseltamivir is temporarily unavailable.

Background

Antiviral treatment of influenza is an important adjunct to influenza vaccination in the prevention and control of influenza and, when given early, reduces the duration of symptoms and may reduce the risk of some complications [2-4]. Influenza viruses typically circulate annually in the United States, most commonly from the late fall through the early spring. Most people recover from influenza without serious complications or sequelae. However, influenza can be associated with serious illnesses, hospitalizations, and deaths, particularly among [people at increased risk](#)



[of complications](#) such as older adults, very young children, pregnant people, and people of all ages with certain chronic medical conditions [2].

Four FDA approved prescription [antiviral medications](#) (oseltamivir, baloxavir, zanamivir, and peramivir) are available for use for early treatment of outpatients with influenza. These antivirals have different formulations, routes of administration, dosing, duration of treatment, and recommendations for administration by age group. The clinical benefit of antiviral treatment of influenza is greatest when treatment is started early (within 2 days of illness onset) in people with mild, uncomplicated illness [3-4]. Oseltamivir treatment also is [recommended](#) as soon as possible for suspected or confirmed influenza requiring hospitalization, and to help control institutional influenza outbreaks [4].

A wide range of [tests](#) for respiratory specimens are available in clinical settings for diagnosing [influenza](#). Use of influenza testing, particularly rapid molecular assays, can inform antiviral treatment decisions, especially when other respiratory viruses are co-circulating in the community. CDC [has testing guidance](#) for clinicians when SARS-CoV-2 and influenza viruses are co-circulating. *Of note, because SARS-CoV-2 and influenza virus co-infection can occur, a positive influenza test result without SARS-CoV-2 testing does not exclude COVID-19, and a positive SARS-CoV-2 test result without influenza testing does not exclude influenza.*

CDC recommends annual [influenza vaccination](#) of people aged 6 months and older as the first and most important tool to prevent influenza [2]. For people aged 65 years and older, high-dose, recombinant, or adjuvanted influenza vaccines are recommended if available [2]. Healthcare providers should strongly encourage people who have not yet received influenza vaccination this season to get vaccinated as they can still benefit.

General Recommendations for Clinicians and Public Health Practitioners

[Available information](#) suggests that current local antiviral availability issues are due to limited availability of *generic* oseltamivir, specifically [1].

- If available, brand-name oseltamivir (Tamiflu) can be used to treat outpatients and hospitalized patients with influenza.
- If oseltamivir is unavailable, [oral baloxavir, inhaled zanamivir, or intravenous peramivir](#) can be used for early treatment of outpatients at increased risk for complications who present with uncomplicated influenza, depending upon age and contraindications.
- When there is limited availability of oseltamivir or other antivirals, antiviral treatment should target patients with influenza who are at the highest risk of severe disease and those who are hospitalized.
- Antiviral treatment of outpatients should be prioritized for persons who test positive for influenza within 2 days of illness onset.
- When there is limited availability of oseltamivir or other antivirals, patients with clinically mild influenza who are otherwise healthy and not at increased risk of influenza complications can be managed with supportive care without antiviral treatment.

Influenza Testing Considerations

- When antivirals are available, a clinical diagnosis of influenza without influenza testing can be made to support prescribing empiric antiviral treatment in outpatients.
- However, in settings where oseltamivir is currently unavailable, influenza testing for patients with suspected influenza is highly recommended to guide antiviral treatment.



- When there are limited supplies of antivirals, treatment of suspected influenza without a positive test result should be limited to those who are being hospitalized with suspected influenza, or patients highly suspected to have influenza (e.g., an ill patient who has a household member with laboratory-confirmed influenza).

Clinicians, hospitals, healthcare systems, and public health officials are encouraged to use all available information and their best judgment to prioritize oseltamivir and other antivirals for treating patients with influenza, depending upon their local situation. The following are considerations for antiviral treatment prioritization when antivirals such as oseltamivir are in short supply.

Guidance for Prioritization when Antiviral Supplies are Limited Hospitalized Patients

- **Prioritize oseltamivir treatment as soon as possible for hospitalized patients with suspected or laboratory-confirmed influenza.**
 - Oseltamivir is the only antiviral that is [recommended](#) for treating influenza in hospitalized patients [4]. Because observational studies have shown that early initiation of oseltamivir treatment has significant clinical benefit [5-7], oseltamivir treatment is recommended to be started as soon as possible without waiting for results of influenza testing, such as in the emergency department or in admitted patients with high suspicion for influenza.
 - There are limited data for using inhaled zanamivir, intravenous peramivir, or baloxavir for treating influenza in hospitalized patients.
 - In a recent clinical trial, the addition of baloxavir to a neuraminidase inhibitor (primarily oseltamivir) did not show clinical benefit compared to neuraminidase inhibitor treatment alone in hospitalized patients with influenza aged 12 years and older [8].

Outpatients

Among outpatients, prioritize antiviral treatment for patients who test positive for influenza as follows:

- **Patients at increased risk of influenza complications and who test positive for influenza within 2 days of illness onset.**
 - People with multiple [conditions](#) that place them at increased risk for complications from influenza (e.g., several co-morbidities, age <2 years, and 65 years and older) and those with severe uncontrolled chronic disease might be at highest risk of influenza complications.
- **Patients who have progressive or severe influenza not requiring hospitalization, even if they test positive for influenza more than 2 days from illness onset.**
- **Patients who are pregnant, less than 2 weeks postpartum, or immunocompromised.**
 - Substantial data from observational studies indicate that oseltamivir treatment of influenza is safe in pregnancy [9].
 - There are no data on the safety or efficacy of baloxavir in pregnancy and baloxavir is [not recommended](#) for pregnant people or those less than 2 weeks postpartum [9].
 - Treatment with a neuraminidase inhibitor (oseltamivir, zanamivir, or peramivir) is recommended for immunocompromised people with influenza.



- Baloxavir is not recommended for treating influenza in immunocompromised people because the optimal duration of treatment is unknown and there is concern for emergence of influenza viruses resistant to baloxavir during or after treatment.
- **Children less than 5 years of age.**
 - Oseltamivir is the only recommended oral antiviral for treatment of influenza in children less than 5 years of age.
 - If oseltamivir suspension is unavailable for treating influenza in young children, clinicians can request that pharmacists compound a suspension from oseltamivir [capsules](#).
 - In areas where generic oseltamivir is unavailable, baloxavir can be used for early treatment of influenza in otherwise healthy children aged 5 years and older, and for children aged 12 years and older with underlying conditions that increase their risk of influenza complications.

Institutional Settings

- **When an influenza outbreak is not occurring, prioritize oseltamivir for early treatment of influenza in residents of congregate settings such as long-term care facilities (LTCFs), who test positive for influenza.**
- **In the setting of laboratory confirmed [influenza outbreaks in LTCFs](#):**
 - Early empiric antiviral treatment of suspected influenza in residents is [recommended](#) [4]. Once an influenza diagnosis is confirmed through testing, post-exposure antiviral chemoprophylaxis of exposed residents is [recommended](#) [4].
 - Because institutional outbreaks can be prolonged, consider using a limited duration treatment dosage (twice daily for 5 days) for post-exposure oseltamivir instead of extended use of oseltamivir chemoprophylaxis (once daily), with ongoing active daily monitoring and influenza testing for all residents with new illness signs and symptoms.
 - If oseltamivir is not available, baloxavir, zanamivir, or peramivir may be used for treatment of influenza.
 - Although baloxavir may be used for treatment, there are no available data on using baloxavir in LTCFs for treatment or post-exposure chemoprophylaxis.

Other Considerations

- In hospitalized patients, oseltamivir can be administered orally or enterically via oro- or nasogastric tube. For hospitalized patients who cannot absorb enterically-administered oseltamivir (e.g., due to gastric stasis, malabsorption, or gastrointestinal bleeding), or when oseltamivir is not available, intravenous peramivir is an option.
- For children who are not able to swallow prescribed oseltamivir capsules, [the prescribed capsules may be opened and mixed with a thick sweetened liquid, such as chocolate syrup, prior to administration](#).
- When local generic oseltamivir availability issues are resolved, CDC recommends reverting back to [original antiviral recommendations](#) that include clinical diagnosis and empiric antiviral treatment of influenza in outpatients.
- Healthcare providers should use clinical judgement and all available data when making decisions about prescribing antibiotics to patients presenting with acute respiratory illness



For More Information

- CDC. [Information for Clinicians on Influenza Virus Testing.](#)
- CDC. [Influenza Antiviral Medications: Summary for Clinicians.](#)
- CDC. [Interim Guidance for Influenza Outbreak Management in Long-Term Care and Post-Acute Care Facilities.](#)
- CDC. [Testing and Management Considerations for Nursing Home Residents with Acute Respiratory Illness Symptoms when SARS-CoV-2 and Influenza Viruses are Co-circulating.](#)

References

1. American Society of Healthcare Pharmacists. Current Drug Shortages. Accessed at: <https://www.ashp.org/drug-shortages/current-shortages>
2. Grohskopf LA, Blanton LH, Ferdinands JM, et al. Prevention and Control of Seasonal Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices — United States, 2022–23 Influenza Season. *MMWR Recomm Rep* 2022;71(No. RR-1):1–28. DOI: <http://dx.doi.org/10.15585/mmwr.r7101a1>
3. Uyeki TM, Hui DS, Zambon M, Wentworth DE, Monto AS. Influenza. *Lancet*. 2022 Aug 27;400(10353):693-706. DOI: [https://doi.org/10.1016/S0140-6736\(22\)00982-5](https://doi.org/10.1016/S0140-6736(22)00982-5)
4. Uyeki TM, Bernstein HH, Bradley JS et al. Clinical Practice Guidelines by the Infectious Diseases Society of America: 2018 Update on Diagnosis, Treatment, Chemoprophylaxis, and Institutional Outbreak Management of Seasonal Influenza. *Clin Infect Dis*. 2019 Mar 5;68(6):e1-e47. DOI: <https://doi.org/10.1093/cid/ciz044>
5. Venkatesan S, Myles PR, Bolton KJ et al. Neuraminidase Inhibitors and Hospital Length of Stay: A Meta-analysis of Individual Participant Data to Determine Treatment Effectiveness Among Patients Hospitalized with Nonfatal 2009 Pandemic Influenza A(H1N1) Virus Infection. *J Infect Dis*. 2020 Jan 14;221(3):356-366. DOI: <https://doi.org/10.1093/infdis/jiz152>
6. Katzen J, Kohn R, Houk JL et al. Early Oseltamivir After Hospital Admission Is Associated with Shortened Hospitalization: A 5-Year Analysis of Oseltamivir Timing and Clinical Outcomes. *Clin Infect Dis*. 2019 Jun 18;69(1):52-58. DOI: <https://doi.org/10.1093/cid/ciy860>
7. Walsh PS, Schnadower D, Zhang Y et al. Association of Early Oseltamivir with Improved Outcomes in Hospitalized Children With Influenza, 2007-2020. *JAMA Pediatr*. 2022 Nov 1;176(11):e223261. DOI: [10.1001/jamapediatrics.2022.3261](https://doi.org/10.1001/jamapediatrics.2022.3261)
8. Kumar D, Ison MG, Mira JP et al. Combining baloxavir marboxil with standard-of-care neuraminidase inhibitor in patients hospitalised with severe influenza (FLAGSTONE): a randomised, parallel-group, double-blind, placebo-controlled, superiority trial. *Lancet Infect Dis*. 2022 May;22(5):718-730. DOI: [https://doi.org/10.1016/S1473-3099\(21\)00469-2](https://doi.org/10.1016/S1473-3099(21)00469-2)
9. Chow EJ, Beigi RH, Riley LE et al. Clinical Effectiveness and Safety of Antivirals for Influenza in Pregnancy. *Open Forum Infect Dis*. 2021 Mar 20;8(6):ofab138. DOI: <https://doi.org/10.1093/ofid/ofab138>

The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on critical health issues; and promotes healthy living through strong partnerships with local, national and international organizations.



NEW YORK CITY DEPARTMENT OF
HEALTH AND MENTAL HYGIENE

Ashwin Vasani, MD, PhD
Commissioner

HAN Message Types

- **Health Alert:** Conveys the highest level of importance; warrants immediate action or attention.
- **Health Advisory:** Provides important information for a specific incident or situation; may not require immediate action.
- **Health Update:** Provides updated information regarding an incident or situation; unlikely to require immediate action.
- **Info Service:** Provides general information that is not necessarily considered to be of an emergent nature.

###

This message was distributed to state and local health officers, state and local epidemiologists, state and local laboratory directors, public information officers, HAN coordinators, and clinician organizations.

###