

HIV EPIDEMIOLOGY & FIELD SERVICES SEMIANNUAL REPORT

Covering January 1, 2010 — December 31, 2010

October 2011

The New York City Department of Health and Mental Hygiene

Vol. 6, No. 2

What's in this report?	
Highlights from 2010	1
History of the HIV/AIDS Epidemic in New York City, 1981 – 2010	1
HIV/AIDS Diagnoses and Deaths, January – December 2010, and PLWHA as of December 31, 2010	2
Which HIV-related Events are Reportable?	2
HIV/AIDS Diagnoses and Deaths, January – December 2010, and PLWHA as of December 31, 2010, by Sex	3
Rates of HIV Diagnoses, PWHA, and Deaths among PWHA by United Hospital Fund (UHF) Neighborhood, New York City 2010	4

Publication schedule: This report reflects events occurring through December 31, 2010, and reported by September 30, 2011, unless otherwise stated. It represents diagnoses made through nine months prior to the publication date. Case reporting is 90% complete by that time. Because cases continue to be reported for many months, the final numbers can be expected to be higher.

To receive this report via e-mail, send an e-mail request to: hivreport@health.nyc.gov

For electronic versions of this and other HIV-related reports, visit: http://www.nyc.gov/html/doh/html/dires/hivepi.shtml

HIV Epidemiology and Field Services Program 42-09 28th Street, CN44, Long Island City, NY 11101 Tel: (212) 442-3388 Fax: (347) 396-7793

On September 1, 2010, New York State HIV testing law was amended. For more information:

http://www.health.state.ny.us/diseases/aids/testing/hiv_testing_law.htm

HIGHLIGHTS FROM 2010

- As of September 30, 2011, 110,736 persons had been diagnosed and reported in New York City and were presumed to be living with HIV/AIDS (PLWHA). This represents an increase of 1.7% from 2009, and 12% from 2006.
- In 2010, there were 3,481 new diagnoses of HIV/AIDS in New York City:
 - 76.6% of new diagnoses were among men, and 23.4% were among women. 48.3% of new diagnoses were among men who have sex with men (MSM), while 21.4% were among persons reporting heterosexual transmission risk. 78.6% of new diagnoses were among blacks and Hispanics. Manhattan and Brooklyn had the highest proportions of new diagnoses. 177 persons (5.1% of cases) were newly diagnosed with HIV in the acute stage of infection.
- □ Comparing 2010 with previous years:
 - New diagnoses declined 8% (from 3,782 to 3,481) between 2009 and 2010; new diagnoses have declined 16% since 2006.
 - Deaths from all causes declined 9% (from 1,871 to 1,695) between 2009 and 2010; deaths from all causes have declined 30% since 2006.
 - The proportion of new HIV cases diagnosed concurrently with AIDS remained stable between 2009 and 2010 (21% in 2009 and 21.8% in 2010).
 - 52.1% of new HIV (non-AIDS) diagnoses in 2010 were among MSM versus 47.4% in 2009.

History of the HIV/AIDS Epidemic in New York City, 1981–2010




Table 1. HIV/AIDS diagnoses and deaths occurring January 1, 2010, through December 31, 2010, and persons diagnosed and reported in New York City and presumed to be living with HIV/AIDS as of December 31, 2010¹

	HIV diagnoses ¹							AIDS diagnoses ³		PLWHA as of 12/31/2010		Deaths ⁴	
	Total		Without AIDS		Concurrent with AIDS diagnosis ²								
	N	%	N	%	N	%	Row %	N	%	N	%	N	%
Total	3,481	100.0	2,722	100.0	759	100.0	21.8	2,520	100.0	110,736	100.0	1,695	100.0
Sex													
Male	2,668	76.6	2,096	77.0	572	75.4	21.4	1,809	71.8	78,663	71.0	1,199	70.7
Female	813	23.4	626	23.0	187	24.6	23.0	711	28.2	32,073	29.0	496	29.3
Race/Ethnicity ⁵													
Black	1,661	47.7	1.273	46.8	388	51.1	23.4	1,322	52.5	49.530	44.7	880	51.9
Hispanic	1,076	30.9	851	31.3	225	29.6	20.9	786	31.2	36,021	32.5	577	34.0
White	638	18.3	516	19.0	122	16.1	19.1	367	14.6	22,656	20.5	220	13.0
Asian/Pacific Islander	95	2.7	72	2.6	23	3.0	24.2	41	1.6	1,808	1.6	13	0.8
Native American	6	0.2	5	0.2	1	0.1	16.7	3	0.1	256	0.2	4	0.2
Multiracial	5	0.1	5	0.2	0	0.0	0.0	1	0.0	50	0.0	1	0.1
Unknown	0	0.0	0	0.0	0	0.0	0.0	0	0.0	415	0.4	0	0.0
Age group (years) ⁶													
0-12	9	0.3	8	0.3	1	0.1	11.1	2	0.1	321	0.3	0	0.0
13-19	135	3.9	114	4.2	21	2.8	15.6	63	2.5	1,336	1.2	4	0.2
20-29	1,076	30.9	951	34.9	125	16.5	11.6	422	16.7	8,051	7.3	26	1.5
30-39	840	24.1	667	24.5	173	22.8	20.6	584	23.2	16,568	15.0	118	7.0
40-49	791	22.7	585	21.5	206	27.1	26.0	770	30.6	38,242	34.5	461	27.2
50-59	447	12.8	284	10.4	163	21.5	36.5	498	19.8	32,153	29.0	650	38.3
60+	183	5.3	113	4.2	70	9.2	38.3	181	7.2	14,065	12.7	436	25.7
Borough of residence													
Bronx	694	19.9	536	19.7	158	20.8	22.8	602	23.9	23,362	21.1	416	24.5
Brooklyn	904	26.0	665	24.4	239	31.5	26.4	760	30.2	27,087	24.5	528	31.2
Manhattan	935	26.9	771	28.3	164	21.6	17.5	543	21.5	33,104	29.9	430	25.4
Queens	565	16.2	429	15.8	136	17.9	24.1	377	15.0	15,824	14.3	195	11.5
Staten Island	57	1.6	40	1.5	17	2.2	29.8	55	2.2	1,900	1.7	48	2.8
Outside NYC	309	8.9	264	9.7	45	5.9	14.6	157	6.2	8,692	7.8	69	4.1
Unknown	17	0.5	17	0.6	0	0.0	0.0	26	1.0	767	0.7	9	0.5
Transmission risk													
Men who have sex with men	1,683	48.3	1,419	52.1	264	34.8	15.7	856	34.0	37,720	34.1	274	16.2
Injection drug use history	164	4.7	128	4.7	36	4.7	22.0	280	11.1	20,505	18.5	636	37.5
Heterosexual ⁷	745	21.4	539	19.8	206	27.1	27.7	612	24.3	21,253	19.2	309	18.2
Perinatal	12	0.3	8	0.3	4	0.5	33.3	40	1.6	2,477	2.2	8	0.5
Other	0	0.0	0	0.0	0	0.0	0.0	0	0.0	247	0.2	9	0.5
Unknown	877	25.2	628	23.1	249	32.8	28.4	732	29.0	28,534	25.8	459	27.1
Clinical status as of 12/31/2010													
HIV (non-AIDS)	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	44,317	40.0	282	16.6
AIDS	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	66,419	60.0	1,413	83.4

PLWHA=Persons living with HIV/AIDS; n/a=Not applicable. All percents are column percents unless otherwise indicated.

Which HIV-related events are reportable in New York State, and who is required to report?

In 1998, New York State expanded AIDS case reporting to include HIV (Chapter 163 of the Laws of 1998, PHL Article 21). The law took effect on **June 1, 2000**, and was expanded further on **June 1, 2005**. All diagnostic and clinical providers (doctors, nurses, physician assistants, and all others diagnosing HIV or providing care to HIV-infected persons) and laboratories are required by law to report the following events:

Events reportable by providers on the required New York State Provider Report Form (PRF)

- Diagnoses of HIV infection
- Diagnoses of HIV illness in a previously unreported individual (i.e., HIV illness not meeting the AIDS case definition)
- Diagnoses of AIDS-defining conditions

Events reportable by laboratories

- All positive Western blot test results
- All viral load test results (detectable and undetectable)
- All CD4 test results
- All viral nucleotide sequence results

For assistance in reporting a case of HIV/AIDS, to receive Provider Report Forms, or to obtain more information, please call (212) 442-3388



TALK TO US CNAP (212) 693-1419

New York State law also requires medical providers to talk to HIV-infected patients about partner notification and report such patients' known sex or needle-sharing partners to the health department. Providers can fill in the *Medical Provider HIV/AIDS and Partner/Contact Report Form* (PRF) or call the NYC DOHMH Contact Notification Assistance Program (CNAP) at (212) 693-1419. Providers can refer HIV-infected patients to CNAP for direct assistance with partner notification.

For more information about the New York State HIV reporting and partner notification law and CNAP, visit: http://www.nyc.gov/html/doh/html/dires/hcpreporting_how.shtml.

¹ For events reported by September 30, 2011. ² HIV diagnosed concurrently with AIDS (within 31 days of HIV diagnosis). Row percent is percent of total HIV diagnoses that were concurrent with AIDS diagnoses. ³ AIDS was diagnosed in 2010 and includes concurrent HIV/AIDS diagnosis. ⁴ Includes deaths from any cause in persons with HIV/AIDS. ⁵ For technical notes on race/ethnicity: http://www.nyc.gov/html/doh/html/ah/hivtables.shtml#abbrev. ⁶ For HIV and AIDS diagnoses, age at diagnosis; for PLWHA, age as of December 31, 2010; and for deaths, age at death. ⁷ Includes persons who had heterosexual sex with a person they know to be HIV-infected, an injection drug user, or a person who has received blood products. For females only, also includes history of prostitution, multiple sex partners, sexually transmitted disease, crack/cocaine use, sex with a bisexual male, probable heterosexual transmission as noted in medical chart, or sex with a male and negative history of injection drug use.


Table 2. HIV/AIDS diagnoses and deaths occurring January 1, 2010, through December 31, 2010, and persons diagnosed and reported in New York City and presumed to be living with HIV/AIDS as of December 31, 2010¹, by sex


	HIV diagnoses ¹					AIDS diagnoses ³		PLWHA as of 12/31/2010		Deaths ⁴			
	т.	otal	Concurrent with Without AIDS AIDS diagnosis ²										
	N	%	N	%	N	%	Row %	N	%	N	%	N	%
Total (Male and Female)	3,481	100.0	2,722	100.0	759	100.0	21.8	2,520	100.0	110,736	100.0	1,695	100.0
Male	2,668	100.0	2,096	100.0	572	100.0	21.4	1,809	100.0	78,663	100.0	1,199	100.0
Race/Ethnicity ⁵ Black	1,144	42.9	884	42.2	260	45.5	22.7	862	47.7	30,860	39.2	596	49.7
Hispanic	843	31.6	665	31.7	178	31.1	21.1	577	31.9	25,593	32.5	421	35.1
White	591	22.2	479	22.9	112	19.6	19.0	330	18.2	20,202	25.7	173	14.4
Asian/Pacific Islander Native American	82 3	3.1 0.1	61 2	2.9 0.1	21 1	3.7 0.2	25.6 33.3	37 2	2.0 0.1	1,470 183	1.9 0.2	7 1	0.6 0.1
Multiracial	5 5	0.1	5	0.1	0	0.2	0.0	1	0.1	37	0.2	1	0.1
Unknown	0	0.0	0	0.0	Ō	0.0	0.0	0	0.0	318	0.4	0	0.0
Age group (years) ⁶	_	0.0	-	0.0	0	0.0	0.0		0.4	450	0.0	0	0.0
0-12 13-19	5 101	0.2 3.8	5 86	0.2 4.1	0 15	0.0 2.6	0.0 14.9	1 40	0.1 2.2	159 669	0.2 0.9	0 3	0.0 0.3
20-29	915	34.3	810	38.6	105	18.4	11.5	345	19.1	5,934	7.5	22	1.8
30-39	658	24.7	523	25.0	135	23.6	20.5	416	23.0	11,523	14.6	80	6.7
40-49 50-59	572 291	21.4 10.9	423 176	20.2 8.4	149 115	26.0 20.1	26.0 39.5	532 344	29.4 19.0	26,874 22,968	34.2 29.2	299 463	24.9 38.6
60+	126	4.7	73	3.5	53	9.3	42.1	131	7.2	10,536	13.4	332	27.7
Borough of residence													
Bronx Brooklyn	465 642	17.4 24.1	361 474	17.2 22.6	104 168	18.2 29.4	22.4 26.2	383 513	21.2 28.4	14,031 17,357	17.8 22.1	297 345	24.8 28.8
Manhattan	808	30.3	665	31.7	143	25.0	17.7	454	25.1	27,017	34.3	333	27.8
Queens	430	16.1	327	15.6	103	18.0	24.0	275	15.2	11,262	14.3	136	11.3
Staten Island Outside NYC	45 263	1.7 9.9	30 224	1.4 10.7	15 39	2.6 6.8	33.3 14.8	42 129	2.3 7.1	1,231 7,196	1.6 9.1	32 50	2.7 4.2
Unknown	15	0.6	15	0.7	0	0.0	0.0	13	0.7	569	0.7	6	0.5
Transmission risk													
Men who have sex with men	1,683	63.1	1,419	67.7	264	46.2	15.7	856	47.3	37,720	48.0	274	22.9
Injection drug use history Heterosexual ⁷	125 145	4.7 5.4	92 93	4.4 4.4	33 52	5.8 9.1	26.4 35.9	200 162	11.1 9.0	14,533 5,669	18.5 7.2	483 98	40.3 8.2
Perinatal	6	0.2	5	0.2	1	0.2	16.7	14	0.8	1,208	1.5	4	0.3
Other	0	0.0	0	0.0	0	0.0	0.0	0	0.0	125	0.2	4	0.3
Unknown Clinical status as of 12/31/2010	709	26.6	487	23.2	222	38.8	31.3	577	31.9	19,408	24.7	336	28.0
HIV (non-AIDS)	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	31,401	39.9	202	16.8
AIDS	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	47,262	60.1	997	83.2
Female	813	100.0	626	100.0	187	100.0	23.0	711	100.0	32,073	100.0	496	100.0
Race/Ethnicity ⁵	F.17	00.0	000	00.4	400	00.4	04.0	400	04.7	40.070	50.0	004	57.0
Black Hispanic	517 233	63.6 28.7	389 186	62.1 29.7	128 47	68.4 25.1	24.8 20.2	460 209	64.7 29.4	18,670 10,428	58.2 32.5	284 156	57.3 31.5
White	47	5.8	37	5.9	10	5.3	21.3	37	5.2	2,454	7.7	47	9.5
Asian/Pacific Islander	13	1.6	11	1.8	2	1.1	15.4	4	0.6	338	1.1	6	1.2
Native American Multiracial	3 0	0.4 0.0	3 0	0.5 0.0	0 0	0.0 0.0	0.0 0.0	1 0	0.1 0.0	73 13	0.2 0.0	3 0	0.6 0.0
Unknown	0	0.0	Ö	0.0	Ö	0.0	0.0	Ö	0.0	97	0.3	ő	0.0
Age group (years) ⁶										400			
0-12 13-19	4 34	0.5 4.2	3 28	0.5 4.5	1 6	0.5 3.2	25.0 17.6	1 23	0.1 3.2	162 667	0.5 2.1	0 1	0.0 0.2
20-29	161	19.8	141	22.5	20	10.7	12.4	77	10.8	2,117	6.6	4	0.8
30-39	182	22.4	144	23.0	38	20.3	20.9	168	23.6	5,045	15.7	38	7.7
40-49 50-59	219 156	26.9 19.2	162 108	25.9 17.3	57 48	30.5 25.7	26.0 30.8	238 154	33.5 21.7	11,368 9,185	35.4 28.6	162 187	32.7 37.7
60+	57	7.0	40	6.4	17	9.1	29.8	50	7.0	3,529	11.0	104	21.0
Borough of residence								2.42					
Bronx Brooklyn	229 262	28.2 32.2	175 191	28.0 30.5	54 71	28.9 38.0	23.6 27.1	219 247	30.8 34.7	9,331 9,730	29.1 30.3	119 183	24.0 36.9
Manhattan	127	15.6	106	16.9	21	11.2	16.5	89	12.5	6,087	19.0	97	19.6
Queens	135	16.6	102	16.3	33	17.6	24.4	102	14.3	4,562	14.2	59	11.9
Staten Island Outside NYC	12 46	1.5 5.7	10 40	1.6 6.4	2 6	1.1 3.2	16.7 13.0	13 28	1.8 3.9	669 1,496	2.1 4.7	16 19	3.2 3.8
Unknown	2	0.2	2	0.4	0	0.0	0.0	13	3.9 1.8	1,496	0.6	3	3.6 0.6
Transmission risk													
Injection drug use history	39	4.8	36 446	5.8	3 154	1.6	7.7 25.7	80 450	11.3	5,972	18.6	153	30.8
Heterosexual ⁷ Perinatal	600 6	73.8 0.7	446 3	71.2 0.5	154 3	82.4 1.6	25.7 50.0	450 26	63.3 3.7	15,584 1,269	48.6 4.0	211 4	42.5 0.8
Other	0	0.0	0	0.0	0	0.0	0.0	0	0.0	122	0.4	5	1.0
Unknown	168	20.7	141	22.5	27	14.4	16.1	155	21.8	9,126	28.5	123	24.8
Clinical status as of 12/31/2010 HIV (non-AIDS)	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	12,916	40.3	80	16.1
AIDS	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	19,157	59.7	416	83.9


PLWHA=Persons living with HIV/AIDS; n/a=Not applicable. All percents are column percents unless otherwise indicated. Footnotes appear at the bottom of Table 1.

Rates of HIV diagnoses, PWHA, and deaths among PWHA by United Hospital Fund (UHF) neighborhood, New York City 2010¹

	HIV diagnoses ² per 100,000 population	Reported PWHA as percent of population	Age-adjusted death rate per 1,000 PWHA ³	Population from 2010 Census
Total	42.6	1.4	15.1	8,175,133
Bronx ⁴	50.2	1.7	17.9	1,382,480
Crotona – Tremont	62.1	2.2	18.8	206,116
Fordham – Bronx Park	53.8	1.7	14.7	252,655
High Bridge – Morrisania	58.8	2.4	22.5	207,631
Hunts Point – Mott Haven	71.0	2.3	19.5	136,591
Kingsbridge – Riverdale	23.1	0.6	15.9*	90,892
Northeast Bronx	41.4	0.9	17.0	190,668
Pelham – Throgs Neck	36.6	1.2	12.7	297,927
Brooklyn ⁴	36.1	1.1	19.5	2,504,700
Bedford/Stuyvesant - Crown Heights	69.0	2.1	22.6	318,898
Bensonhurst – Bay Ridge	4.0*	0.3	26.5	199,271
Borough Park	9.9	0.3	11.4	331,983
Canarsie – Flatlands	34.4	0.7	14.7	195,027
Coney Island - Sheepshead Bay	13.0	0.4	22.4	285,502
Downtown - Heights - Park Slope	37.5	1.4	12.6	224,199
East Flatbush – Flatbush	59.3	1.5	15.6	296,583
East New York	55.4	1.4	23.7	187,855
Greenpoint	18.1	8.0	23.6	127,051
Sunset Park	25.8	0.6	11.5*	127,863
Williamsburg – Bushwick	56.1	1.7	24.7	210,468
Manhattan ⁴	59.3	2.1	12.3	1,577,279
Central Harlem – Morningside Heights	98.4	2.8	17.8	162,652
Chelsea - Clinton	140.1	4.3	7.2	144,896
East Harlem	77.3	2.8	18.2	109,972
Gramercy Park – Murray Hill	47.6	1.7	9.2	134,520
Greenwich Village – SoHo	66.9	2.7	11.0	83,749
Lower Manhattan	45.1	0.9	5.7*	53,159
Union Square – Lower East Side	38.7	1.7	14.3	198,781
Upper East Side	15.4	0.7	10.1	220,962
Upper West Side	30.9	1.5	12.6	220,080
Washington Heights – Inwood	58.8	1.7	13.7	248,508
Queens ⁴	25.3	0.7	12.7	2,235,260
Bayside – Little Neck	4.5*	0.2	9.9*	87,972
Flushing – Clearview	4.6	0.3	11.3*	259,767
Fresh Meadows	8.3*	0.3	0.0*	96,831
Jamaica	41.1	1.0	19.3	289,314
Long Island City – Astoria	39.6	0.9	11.4	204,715
Ridgewood – Forest Hills	12.6	0.4	12.4	245,746
Rockaway	36.5	8.0	17.8	114,978
Southeast Queens	21.7	0.6	13.6	189,171
Southwest Queens	22.2	0.5	9.7	266,265
West Queens	34.1	1.0	9.3	480,501
Staten Island ⁴	12.2	0.4	25.0	468,730
Port Richmond	12.8*	0.6	16.3*	70,387
South Beach - Tottenville	8.5	0.2	11.6*	189,185
Stapleton – St. George	23.5	0.7	32.5	123,648
Willowbrook	3.5*	0.2	25.8*	85,510







PWHA=Persons with HIV/AIDS alive at any time January 1, 2010 – December 31, 2010. Map categories defined by neighborhood quartile. Rates based on numerators ≤10 are marked with an asterisk (*) and should be interpreted with caution.

¹ Based on data reported through September 30, 2011.

² Includes diagnoses of HIV without AIDS and HIV concurrent with AIDS.

³ Age-adjusted to the citywide population of PWHA in 2010.

⁴ Borough-level data include persons residing in unknown UHF neighborhoods.